



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:01 AM JST

PDB ID : 9UUV / pdb_00009uuv
Title : beta barrel protein-Luck
Authors : Zhang, B.; Ge, H.M.; Ma, X.X.
Deposited on : 2025-05-08
Resolution : 1.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.48.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1090 (1.88-1.88)
Clashscore	180529	1144 (1.88-1.88)
Ramachandran outliers	177936	1135 (1.88-1.88)
Sidechain outliers	177891	1135 (1.88-1.88)
RSRZ outliers	164620	1090 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	172	3% 70%12%•17%
1	B	172	4% 71%10%•18%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4214 atoms, of which 1880 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Allene oxide cyclase barrel-like domain-containing protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	142	Total	C	H	N	O	S	0	0	0
			2034	701	940	181	210	2			
1	B	141	Total	C	H	N	O	S	0	0	0
			2026	695	940	179	211	1			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP A0AAU3BQU8
A	-18	GLY	-	expression tag	UNP A0AAU3BQU8
A	-17	SER	-	expression tag	UNP A0AAU3BQU8
A	-16	SER	-	expression tag	UNP A0AAU3BQU8
A	-15	HIS	-	expression tag	UNP A0AAU3BQU8
A	-14	HIS	-	expression tag	UNP A0AAU3BQU8
A	-13	HIS	-	expression tag	UNP A0AAU3BQU8
A	-12	HIS	-	expression tag	UNP A0AAU3BQU8
A	-11	HIS	-	expression tag	UNP A0AAU3BQU8
A	-10	HIS	-	expression tag	UNP A0AAU3BQU8
A	-9	SER	-	expression tag	UNP A0AAU3BQU8
A	-8	SER	-	expression tag	UNP A0AAU3BQU8
A	-7	GLY	-	expression tag	UNP A0AAU3BQU8
A	-6	LEU	-	expression tag	UNP A0AAU3BQU8
A	-5	VAL	-	expression tag	UNP A0AAU3BQU8
A	-4	PRO	-	expression tag	UNP A0AAU3BQU8
A	-3	ARG	-	expression tag	UNP A0AAU3BQU8
A	-2	GLY	-	expression tag	UNP A0AAU3BQU8
A	-1	SER	-	expression tag	UNP A0AAU3BQU8
A	0	HIS	-	expression tag	UNP A0AAU3BQU8
B	-19	MET	-	initiating methionine	UNP A0AAU3BQU8
B	-18	GLY	-	expression tag	UNP A0AAU3BQU8
B	-17	SER	-	expression tag	UNP A0AAU3BQU8
B	-16	SER	-	expression tag	UNP A0AAU3BQU8
B	-15	HIS	-	expression tag	UNP A0AAU3BQU8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP A0AAU3BQU8
B	-13	HIS	-	expression tag	UNP A0AAU3BQU8
B	-12	HIS	-	expression tag	UNP A0AAU3BQU8
B	-11	HIS	-	expression tag	UNP A0AAU3BQU8
B	-10	HIS	-	expression tag	UNP A0AAU3BQU8
B	-9	SER	-	expression tag	UNP A0AAU3BQU8
B	-8	SER	-	expression tag	UNP A0AAU3BQU8
B	-7	GLY	-	expression tag	UNP A0AAU3BQU8
B	-6	LEU	-	expression tag	UNP A0AAU3BQU8
B	-5	VAL	-	expression tag	UNP A0AAU3BQU8
B	-4	PRO	-	expression tag	UNP A0AAU3BQU8
B	-3	ARG	-	expression tag	UNP A0AAU3BQU8
B	-2	GLY	-	expression tag	UNP A0AAU3BQU8
B	-1	SER	-	expression tag	UNP A0AAU3BQU8
B	0	HIS	-	expression tag	UNP A0AAU3BQU8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	77	Total O 77 77	0	0
2	B	77	Total O 77 77	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Allene oxide cyclase barrel-like domain-containing protein



- Molecule 1: Allene oxide cyclase barrel-like domain-containing protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.40Å 83.07Å 88.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.94 – 1.87 31.94 – 1.87	Depositor EDS
% Data completeness (in resolution range)	99.5 (31.94-1.87) 99.5 (31.94-1.87)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 1.87Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.193 , 0.224 0.193 , 0.225	Depositor DCC
R_{free} test set	1395 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	32.4	Xtriage
Anisotropy	0.251	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4214	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/1114	0.63	0/1508
1	B	0.48	0/1107	0.65	0/1498
All	All	0.50	0/2221	0.64	0/3006

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	37	TYR	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1094	940	1088	24	0
1	B	1086	940	1064	14	0
2	A	77	0	0	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	77	0	0	3	1
All	All	2334	1880	2152	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:26:SER:O	2:B:201:HOH:O	1.86	0.92
1:A:26:SER:O	1:A:41:LYS:NZ	2.08	0.87
1:B:5:THR:HG22	1:B:6:LYS:HD3	1.57	0.87
1:A:151:GLU:O	2:A:201:HOH:O	1.97	0.81
1:A:39:SER:OG	2:A:202:HOH:O	1.99	0.79
1:A:14:LEU:HB2	1:A:142:VAL:CG2	2.16	0.76
1:B:11:ILE:HD11	1:B:122:TYR:OH	1.90	0.72
1:A:39:SER:C	2:A:202:HOH:O	2.32	0.71
1:B:48:TYR:OH	1:B:85:GLU:OE2	2.05	0.68
1:A:11:ILE:HD11	1:A:122:TYR:OH	1.97	0.64
1:B:5:THR:C	2:B:202:HOH:O	2.41	0.63
1:A:100:LEU:HD13	1:A:101:ASP:C	2.24	0.62
1:A:110:ILE:HD11	1:A:132:ARG:NH2	2.15	0.62
1:B:5:THR:N	2:B:202:HOH:O	2.31	0.62
1:B:35:GLY:O	1:B:36:ASP:HB3	2.00	0.61
1:A:14:LEU:O	1:A:142:VAL:HG22	2.03	0.59
1:A:100:LEU:HD22	1:A:101:ASP:H	1.67	0.58
1:A:14:LEU:C	1:A:142:VAL:HG22	2.29	0.57
1:A:110:ILE:HD11	1:A:132:ARG:HH22	1.69	0.57
1:A:74:GLU:H	1:A:74:GLU:CD	2.13	0.57
1:B:151:GLU:O	1:B:152:ALA:HB3	2.07	0.54
1:A:39:SER:CA	2:A:202:HOH:O	2.58	0.51
1:A:110:ILE:N	1:A:110:ILE:HD12	2.26	0.50
1:A:110:ILE:HD12	1:A:110:ILE:H	1.76	0.50
1:A:1:MET:N	2:A:203:HOH:O	2.21	0.49
1:A:100:LEU:HD22	1:A:101:ASP:N	2.27	0.49
1:A:39:SER:O	2:A:202:HOH:O	2.20	0.49
1:A:84:ARG:NH1	1:A:86:GLU:OE2	2.29	0.48
1:B:85:GLU:OE1	1:B:129:ARG:NH2	2.47	0.47
1:A:74:GLU:CD	1:A:74:GLU:N	2.73	0.46
1:B:151:GLU:CD	1:B:151:GLU:H	2.24	0.46
1:B:117:GLY:HA3	1:B:122:TYR:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:130:GLU:O	1:B:142:VAL:HA	2.18	0.43
1:B:5:THR:HG22	1:B:6:LYS:CD	2.38	0.42
1:A:142:VAL:O	1:A:142:VAL:HG23	2.18	0.42
1:B:21:LEU:CD2	1:B:23:MET:HE2	2.49	0.42
1:A:135:ASN:OD1	1:A:137:LYS:HG3	2.19	0.41
1:A:81:SER:O	1:A:99:PRO:HA	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:211:HOH:O	2:B:256:HOH:O[1_655]	1.88	0.32

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/172 (80%)	137 (99%)	1 (1%)	0	100	100
1	B	137/172 (80%)	136 (99%)	1 (1%)	0	100	100
All	All	275/344 (80%)	273 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/138 (83%)	110 (96%)	4 (4%)	31	15
1	B	112/138 (81%)	109 (97%)	3 (3%)	40	24
All	All	226/276 (82%)	219 (97%)	7 (3%)	35	18

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	40	VAL
1	A	58	GLN
1	A	74	GLU
1	B	107	GLN
1	B	142	VAL
1	B	151	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	104	GLN
1	B	51	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	142/172 (82%)	0.03	5 (3%) 47 50	26, 40, 60, 85	0
1	B	141/172 (81%)	0.17	7 (4%) 35 36	26, 40, 66, 115	0
All	All	283/344 (82%)	0.10	12 (4%) 41 42	26, 40, 63, 115	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	152	ALA	5.4
1	A	27	VAL	4.7
1	B	5	THR	4.7
1	B	35	GLY	4.3
1	A	152	ALA	3.3
1	B	34	GLY	3.3
1	B	37	TYR	3.1
1	A	38	ASN	2.7
1	A	1	MET	2.7
1	B	6	LYS	2.3
1	B	7	ASP	2.1
1	A	74	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.