



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 21, 2025 – 11:17 PM JST

PDB ID : 9UZE / pdb_00009uze
Title : Crystal structure of Kasokero virus cap- snatching endonuclease (E682A mutant)
Authors : Deng, Z.; Kuang, W.; Tian, Z.
Deposited on : 2025-05-16
Resolution : 2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5670 (2.44-2.40)
Clashscore	180529	6299 (2.44-2.40)
Ramachandran outliers	177936	6232 (2.44-2.40)
Sidechain outliers	177891	6233 (2.44-2.40)
RSRZ outliers	164620	5670 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	63% 16% 20%
2	H	230	85% 10% .
3	L	219	84% 15%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5414 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-directed RNA polymerase L.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	239	Total	C	N	O	S	0	0	0
			1915	1222	317	367	9			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	605	GLY	-	expression tag	UNP A0A0M5KLS1
A	606	PRO	-	expression tag	UNP A0A0M5KLS1
A	607	LEU	-	expression tag	UNP A0A0M5KLS1
A	608	GLY	-	expression tag	UNP A0A0M5KLS1
A	609	SER	-	expression tag	UNP A0A0M5KLS1
A	682	ALA	GLU	engineered mutation	UNP A0A0M5KLS1

- Molecule 2 is a protein called mAb 2E9 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	220	Total	C	N	O	S	0	0	0
			1645	1042	274	322	7			

- Molecule 3 is a protein called mAb 2E9 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	218	Total	C	N	O	S	0	0	0
			1677	1052	286	335	4			

- Molecule 4 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	44	Total 44	O 44	0	0
5	H	69	Total 69	O 69	0	0
5	L	63	Total 63	O 63	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: RNA-directed RNA polymerase L



• Molecule 2: mAb 2E9 Fab heavy chain



• Molecule 3: mAb 2E9 Fab light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	37.18Å 81.11Å 115.87Å 90.00° 93.38° 90.00°	Depositor
Resolution (Å)	29.85 – 2.41 29.85 – 2.41	Depositor EDS
% Data completeness (in resolution range)	92.3 (29.85-2.41) 92.2 (29.85-2.41)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.27 (at 2.42Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.178 , 0.233 0.178 , 0.233	Depositor DCC
R_{free} test set	1265 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å ²)	32.6	Xtriage
Anisotropy	0.203	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5414	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/1957	0.51	0/2650
2	H	0.41	0/1687	0.62	1/2296 (0.0%)
3	L	0.37	0/1714	0.58	0/2324
All	All	0.37	0/5358	0.57	1/7270 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	154	PRO	CA-N-CD	-6.07	103.51	112.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	153	GLU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1867	40	0
2	H	1645	0	1607	16	0
3	L	1677	0	1637	24	1
4	A	1	0	0	0	0
5	A	44	0	0	6	0
5	H	69	0	0	1	0
5	L	63	0	0	1	0
All	All	5414	0	5111	80	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:200:GLU:OE2	5:L:301:HOH:O	1.62	1.15
1:A:617:CYS:O	5:A:1101:HOH:O	1.90	0.89
1:A:661:ARG:NH2	5:A:1102:HOH:O	2.10	0.84
1:A:902:GLU:O	1:A:903:ASN:ND2	2.13	0.80
3:L:185:THR:C	3:L:186:LEU:HD12	2.13	0.74
1:A:826:GLU:HB2	1:A:862:LYS:HG2	1.70	0.73
2:H:131:PRO:HG3	2:H:143:LEU:HB3	1.70	0.72
1:A:722:MET:HE2	1:A:828:LEU:HD23	1.72	0.70
1:A:668:GLU:OE2	1:A:685:LYS:NZ	2.26	0.68
2:H:163:ALA:O	5:H:301:HOH:O	2.10	0.68
1:A:784:GLU:OE2	5:A:1103:HOH:O	2.13	0.67
1:A:835:GLN:HE21	1:A:841:LYS:HG3	1.60	0.65
1:A:615:LEU:O	1:A:619:GLU:HG3	1.97	0.64
1:A:856:MET:HA	1:A:856:MET:HE2	1.79	0.64
1:A:725:SER:HA	1:A:784:GLU:OE1	1.99	0.62
1:A:693:GLN:HE21	1:A:694:THR:HG22	1.64	0.62
3:L:184:LEU:HG	3:L:186:LEU:CD1	2.32	0.60
1:A:809:GLU:HG2	1:A:853:VAL:HG11	1.84	0.59
3:L:186:LEU:HD12	3:L:186:LEU:N	2.19	0.58
2:H:146:LEU:HG	2:H:148:LYS:HG3	1.85	0.57
1:A:835:GLN:NE2	5:A:1104:HOH:O	2.27	0.57
1:A:835:GLN:NE2	1:A:841:LYS:HG3	2.19	0.57
2:H:34:MET:HE1	2:H:98:THR:HB	1.88	0.56
3:L:8:PRO:HG3	3:L:11:LEU:HD13	1.89	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:184:LEU:HG	3:L:186:LEU:HD11	1.89	0.55
2:H:34:MET:CE	2:H:98:THR:HB	2.37	0.55
2:H:34:MET:HB3	2:H:79:LEU:HD22	1.88	0.54
1:A:856:MET:HE2	1:A:859:LEU:HD12	1.90	0.54
2:H:18:ARG:HB3	2:H:83:MET:CE	2.37	0.54
3:L:66:ARG:NH2	3:L:87:ASP:OD1	2.34	0.54
3:L:171:GLN:HG3	3:L:178:TYR:CZ	2.44	0.52
1:A:702:LYS:O	1:A:706:ARG:HG2	2.10	0.52
1:A:856:MET:CE	1:A:859:LEU:HD12	2.40	0.52
1:A:706:ARG:HB2	1:A:855:LEU:HD11	1.93	0.51
1:A:701:GLU:O	1:A:705:GLU:HG3	2.11	0.50
1:A:652:HIS:HD1	1:A:652:HIS:C	2.18	0.50
3:L:186:LEU:CD1	3:L:186:LEU:N	2.73	0.50
3:L:38:LEU:HG	3:L:39:HIS:N	2.27	0.49
1:A:902:GLU:C	1:A:903:ASN:ND2	2.71	0.48
1:A:720:TYR:HB2	1:A:830:VAL:HB	1.95	0.48
3:L:59:ARG:HG2	3:L:63:VAL:HB	1.96	0.48
2:H:93:MET:HE1	2:H:113:LEU:HB2	1.95	0.48
3:L:38:LEU:HD22	3:L:76:PHE:CG	2.49	0.48
1:A:680:ILE:O	1:A:685:LYS:HE2	2.15	0.46
1:A:652:HIS:O	1:A:652:HIS:ND1	2.44	0.46
1:A:901:LEU:HD12	1:A:901:LEU:HA	1.75	0.46
3:L:2:ILE:CD1	3:L:27:GLN:HB2	2.46	0.46
2:H:117:SER:HB3	2:H:151:PHE:HZ	1.80	0.46
1:A:677:TRP:CE2	1:A:722:MET:HE1	2.50	0.45
3:L:125:PRO:HD3	3:L:137:VAL:HG22	1.99	0.45
3:L:1:GLN:OE1	3:L:1:GLN:N	2.33	0.45
3:L:82:ARG:HG2	3:L:82:ARG:HH11	1.81	0.45
1:A:693:GLN:NE2	1:A:694:THR:HG22	2.29	0.44
1:A:827:LEU:HB2	1:A:863:VAL:HG12	1.99	0.44
3:L:37:TYR:O	3:L:95:GLN:HA	2.17	0.44
3:L:23:CYS:HB2	3:L:40:TRP:CH2	2.52	0.44
1:A:892:SER:OG	1:A:893:ILE:N	2.50	0.44
2:H:76:LYS:HB3	2:H:76:LYS:HE3	1.73	0.43
2:H:89:GLU:H	2:H:89:GLU:CD	2.25	0.43
1:A:652:HIS:C	1:A:652:HIS:ND1	2.77	0.43
3:L:11:LEU:HD12	3:L:11:LEU:HA	1.76	0.43
1:A:612:PRO:O	1:A:616:ILE:HG13	2.19	0.43
1:A:898:LYS:O	1:A:902:GLU:HG3	2.19	0.43
1:A:795:PHE:CD2	1:A:866:LEU:HD13	2.54	0.42
1:A:871:ASN:HB2	5:A:1107:HOH:O	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:82:ARG:HG2	3:L:82:ARG:NH1	2.35	0.42
3:L:42:LEU:HD13	3:L:91:TYR:CZ	2.55	0.42
2:H:211:LYS:HZ2	2:H:211:LYS:HG3	1.63	0.42
2:H:110:GLN:OE1	2:H:110:GLN:HA	2.20	0.41
3:L:174:LYS:HA	3:L:174:LYS:HD3	1.85	0.41
3:L:2:ILE:HD13	3:L:27:GLN:HB2	2.01	0.41
2:H:73:ASP:OD1	2:H:75:PRO:HD2	2.21	0.41
2:H:12:VAL:HG22	2:H:18:ARG:HD3	2.02	0.41
1:A:841:LYS:NZ	5:A:1104:HOH:O	2.20	0.41
3:L:139:CYS:HB2	3:L:153:TRP:CH2	2.55	0.41
2:H:34:MET:SD	2:H:98:THR:HB	2.61	0.41
1:A:882:ASP:HB2	1:A:885:TYR:CD2	2.56	0.40
1:A:896:LEU:HG	1:A:900:LEU:HD11	2.02	0.40
1:A:901:LEU:C	1:A:903:ASN:H	2.29	0.40
1:A:823:GLU:HA	1:A:823:GLU:OE1	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:1:GLN:NE2	3:L:84:GLU:OE2[1_655]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/300 (78%)	230 (98%)	5 (2%)	0	100	100
2	H	218/230 (95%)	213 (98%)	4 (2%)	1 (0%)	25	36
3	L	216/219 (99%)	210 (97%)	6 (3%)	0	100	100
All	All	669/749 (89%)	653 (98%)	15 (2%)	1 (0%)	48	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	154	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/271 (78%)	208 (99%)	3 (1%)	62	79
2	H	181/192 (94%)	180 (99%)	1 (1%)	84	92
3	L	193/195 (99%)	191 (99%)	2 (1%)	73	85
All	All	585/658 (89%)	579 (99%)	6 (1%)	73	85

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	836	THR
1	A	894	SER
1	A	901	LEU
2	H	196	THR
3	L	94	SER
3	L	134	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	693	GLN
1	A	783	GLN
1	A	835	GLN
1	A	903	ASN
2	H	110	GLN
2	H	197	GLN
3	L	58	ASN
3	L	165	GLN
3	L	215	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	239/300 (79%)	-0.03	0 100 100	25, 46, 64, 73	0
2	H	220/230 (95%)	-0.54	1 (0%) 87 86	19, 28, 42, 64	0
3	L	218/219 (99%)	-0.57	0 100 100	19, 29, 44, 60	0
All	All	677/749 (90%)	-0.37	1 (0%) 92 92	19, 34, 57, 73	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	154	PRO	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MN	A	1001	1/1	0.98	0.03	56,56,56,56	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.