



Full wwPDB X-ray Structure Validation Report i

Jan 8, 2026 – 12:11 PM JST

PDB ID : 9VC7 / pdb_00009vc7
Title : Structural insights into the tumor suppressor ZMYND11 reveal diverse recognition mechanisms
Authors : Bai, X.; Chen, Z.
Deposited on : 2025-06-05
Resolution : 3.35 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1012 (3.40-3.32)
Clashscore	180529	1035 (3.40-3.32)
Ramachandran outliers	177936	1037 (3.40-3.32)
Sidechain outliers	177891	1037 (3.40-3.32)
RSRZ outliers	164620	1012 (3.40-3.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 7432 atoms, of which 3663 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Zinc finger MYND domain-containing protein 11.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	111	Total	C	H	N	O	S	0	0	0
			1817	563	895	171	175	13			
1	B	112	Total	C	H	N	O	S	7	0	0
			1828	566	901	175	173	13			
1	C	114	Total	C	H	N	O	S	1	0	0
			1904	588	943	183	177	13			
1	D	114	Total	C	H	N	O	S	1	0	0
			1875	582	924	178	179	12			


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		
2	C	2	Total	Zn	0	0
			2	2		
2	D	2	Total	Zn	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Zinc finger MYND domain-containing protein 11




- Molecule 1: Zinc finger MYND domain-containing protein 11



- Molecule 1: Zinc finger MYND domain-containing protein 11



- Molecule 1: Zinc finger MYND domain-containing protein 11



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.14Å 75.95Å 172.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.08 – 3.35 37.08 – 3.35	Depositor EDS
% Data completeness (in resolution range)	98.0 (37.08-3.35) 98.6 (37.08-3.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.96 (at 3.33Å)	Xtriage
Refinement program	PHENIX 2.0_5761	Depositor
R , R_{free}	0.244, 0.262 0.256, 0.273	Depositor DCC
R_{free} test set	649 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å ²)	104.6	Xtriage
Anisotropy	0.503	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35, 76.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7432	wwPDB-VP
Average B, all atoms (Å ²)	130.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/935	0.57	0/1244
1	B	0.29	0/939	0.56	0/1247
1	C	0.17	0/975	0.38	0/1294
1	D	0.17	0/965	0.34	0/1284
All	All	0.23	0/3814	0.47	0/5069

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	526	ARG	Sidechain
1	B	526	ARG	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	922	895	893	12	0
1	B	927	901	896	10	0
1	C	961	943	942	6	0
1	D	951	924	922	9	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
All	All	3769	3663	3653	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:554:ILE:HD13	1:B:554:ILE:HD13	1.79	0.65
1:A:505:MET:HE1	1:B:502:ARG:HA	1.80	0.63
1:D:575:CYS:SG	1:D:594:HIS:CE1	2.93	0.61
1:B:578:THR:HG21	1:B:594:HIS:CE1	2.37	0.60
1:C:550:HIS:HB2	1:D:550:HIS:CE1	2.39	0.57
1:B:570:ALA:HB1	1:B:580:TYR:O	2.05	0.56
1:C:505:MET:HE1	1:D:501:LEU:HD22	1.88	0.55
1:A:523:GLU:HG3	1:B:524:MET:HE1	1.89	0.53
1:C:589:HIS:ND1	1:C:594:HIS:HB2	2.25	0.52
1:D:536:LYS:O	1:D:540:VAL:HG23	2.08	0.52
1:C:539:PHE:HD1	1:D:539:PHE:HD1	1.60	0.48
1:D:590:TRP:HA	1:D:594:HIS:HB3	1.96	0.47
1:A:539:PHE:O	1:A:543:ILE:HG23	2.16	0.46
1:D:571:MET:HE2	1:D:581:CYS:O	2.15	0.46
1:D:501:LEU:O	1:D:501:LEU:HD23	2.16	0.46
1:A:525:ASP:C	1:A:526:ARG:HH21	2.25	0.45
1:A:575:CYS:SG	1:A:594:HIS:CE1	3.10	0.45
1:B:563:CYS:N	1:B:570:ALA:HB2	2.32	0.45
1:A:495:ARG:HG2	1:A:495:ARG:HH11	1.82	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:598:CYS:SG	1:A:599:ARG:N	2.90	0.44
1:C:520:MET:HG2	1:D:520:MET:HB3	2.00	0.44
1:A:579:SER:HG	1:B:576:TRP:HZ3	1.59	0.43
1:A:523:GLU:CG	1:B:524:MET:HE1	2.48	0.43
1:B:527:LYS:O	1:B:531:VAL:HG23	2.19	0.42
1:A:579:SER:OG	1:B:576:TRP:CZ3	2.71	0.41
1:A:527:LYS:O	1:A:531:VAL:HG23	2.21	0.40
1:C:497:ALA:O	1:C:501:LEU:HD23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	109/122 (89%)	101 (93%)	8 (7%)	0	100 100
1	B	108/122 (88%)	103 (95%)	5 (5%)	0	100 100
1	C	112/122 (92%)	104 (93%)	8 (7%)	0	100 100
1	D	112/122 (92%)	108 (96%)	4 (4%)	0	100 100
All	All	441/488 (90%)	416 (94%)	25 (6%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	101/114 (89%)	101 (100%)	0	100 100
1	B	100/114 (88%)	96 (96%)	4 (4%)	27 52
1	C	104/114 (91%)	104 (100%)	0	100 100
1	D	103/114 (90%)	102 (99%)	1 (1%)	73 84
All	All	408/456 (90%)	403 (99%)	5 (1%)	67 80

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	523	GLU
1	B	529	LYS
1	B	530	GLN
1	B	575	CYS
1	D	539	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	530	GLN
1	A	550	HIS
1	A	561	GLN
1	A	591	HIS
1	B	561	GLN
1	C	591	HIS
1	D	550	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	111/122 (90%)	0.17	4 (3%) 46 38	92, 131, 187, 205	0
1	B	112/122 (91%)	0.07	3 (2%) 56 48	69, 128, 159, 200	1 (0%)
1	C	114/122 (93%)	-0.14	0 100 100	87, 120, 156, 200	0
1	D	114/122 (93%)	-0.06	2 (1%) 67 58	94, 130, 164, 224	1 (0%)
All	All	451/488 (92%)	0.01	9 (1%) 64 54	69, 128, 169, 224	2 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	572	TYR	4.6
1	A	586	GLN	3.5
1	B	491	GLU	3.0
1	A	587	GLN	2.9
1	D	488	ARG	2.4
1	A	583	ILE	2.3
1	A	572	TYR	2.3
1	D	572	TYR	2.1
1	B	489	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)


There are no oligosaccharides in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	C	702	1/1	0.98	0.04	120,120,120,120	0
2	ZN	B	701	1/1	0.99	0.03	107,107,107,107	0
2	ZN	B	702	1/1	0.99	0.06	107,107,107,107	0
2	ZN	C	701	1/1	0.99	0.03	100,100,100,100	0
2	ZN	A	702	1/1	0.99	0.03	107,107,107,107	0
2	ZN	D	701	1/1	0.99	0.05	133,133,133,133	0
2	ZN	D	702	1/1	0.99	0.04	111,111,111,111	0
2	ZN	A	701	1/1	1.00	0.04	116,116,116,116	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




















6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.