



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 11, 2026 – 03:17 PM JST

PDB ID : 9VMQ / pdb_00009vmq
Title : Crystal structure of Dot/Icm T4SS effector Ceg10(C159S)
Authors : Ge, H.
Deposited on : 2025-06-29
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.48


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	2% 86% 10% .
1	B	231	2% 87% 9% 5%
1	C	231	% 87% 8% .
1	D	231	2% 86% 9% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7529 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dot/Icm T4SS effector Ceg10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	222	Total	C	N	O	S	0	1	0
			1790	1142	305	338	5			
1	B	220	Total	C	N	O	S	0	0	0
			1780	1135	306	334	5			
1	C	221	Total	C	N	O	S	0	0	0
			1786	1138	307	336	5			
1	D	221	Total	C	N	O	S	0	0	0
			1788	1140	307	336	5			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	55	MET	-	initiating methionine	UNP A0AAN5PG55
A	56	GLY	-	expression tag	UNP A0AAN5PG55
A	57	HIS	-	expression tag	UNP A0AAN5PG55
A	58	HIS	-	expression tag	UNP A0AAN5PG55
A	59	HIS	-	expression tag	UNP A0AAN5PG55
A	60	HIS	-	expression tag	UNP A0AAN5PG55
A	61	HIS	-	expression tag	UNP A0AAN5PG55
A	62	HIS	-	expression tag	UNP A0AAN5PG55
A	63	GLY	-	expression tag	UNP A0AAN5PG55
A	64	SER	-	expression tag	UNP A0AAN5PG55
A	159	SER	CYS	engineered mutation	UNP A0AAN5PG55
B	55	MET	-	initiating methionine	UNP A0AAN5PG55
B	56	GLY	-	expression tag	UNP A0AAN5PG55
B	57	HIS	-	expression tag	UNP A0AAN5PG55
B	58	HIS	-	expression tag	UNP A0AAN5PG55
B	59	HIS	-	expression tag	UNP A0AAN5PG55
B	60	HIS	-	expression tag	UNP A0AAN5PG55
B	61	HIS	-	expression tag	UNP A0AAN5PG55
B	62	HIS	-	expression tag	UNP A0AAN5PG55
B	63	GLY	-	expression tag	UNP A0AAN5PG55
B	64	SER	-	expression tag	UNP A0AAN5PG55

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	159	SER	CYS	engineered mutation	UNP A0AAN5PG55
C	55	MET	-	initiating methionine	UNP A0AAN5PG55
C	56	GLY	-	expression tag	UNP A0AAN5PG55
C	57	HIS	-	expression tag	UNP A0AAN5PG55
C	58	HIS	-	expression tag	UNP A0AAN5PG55
C	59	HIS	-	expression tag	UNP A0AAN5PG55
C	60	HIS	-	expression tag	UNP A0AAN5PG55
C	61	HIS	-	expression tag	UNP A0AAN5PG55
C	62	HIS	-	expression tag	UNP A0AAN5PG55
C	63	GLY	-	expression tag	UNP A0AAN5PG55
C	64	SER	-	expression tag	UNP A0AAN5PG55
C	159	SER	CYS	engineered mutation	UNP A0AAN5PG55
D	55	MET	-	initiating methionine	UNP A0AAN5PG55
D	56	GLY	-	expression tag	UNP A0AAN5PG55
D	57	HIS	-	expression tag	UNP A0AAN5PG55
D	58	HIS	-	expression tag	UNP A0AAN5PG55
D	59	HIS	-	expression tag	UNP A0AAN5PG55
D	60	HIS	-	expression tag	UNP A0AAN5PG55
D	61	HIS	-	expression tag	UNP A0AAN5PG55
D	62	HIS	-	expression tag	UNP A0AAN5PG55
D	63	GLY	-	expression tag	UNP A0AAN5PG55
D	64	SER	-	expression tag	UNP A0AAN5PG55
D	159	SER	CYS	engineered mutation	UNP A0AAN5PG55

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	109	Total O 109 109	0	0
2	B	104	Total O 104 104	0	0
2	C	80	Total O 80 80	0	0
2	D	92	Total O 92 92	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	50.89Å 54.60Å 82.94Å 91.40° 91.11° 91.93°	Depositor
Resolution (Å)	20.00 – 1.79 20.00 – 1.79	Depositor EDS
% Data completeness (in resolution range)	95.8 (20.00-1.79) 95.9 (20.00-1.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 1.78Å)	Xtriage
Refinement program	REFMAC 5.8.0415	Depositor
R, R_{free}	0.154 , 0.199 0.165 , 0.204	Depositor DCC
R_{free} test set	3855 reflections (4.56%)	wwPDB-VP
Wilson B-factor (Å ²)	26.1	Xtriage
Anisotropy	0.133	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 50.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.017 for h,-k,-l 0.220 for -h,k,-l 0.014 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7529	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/1830	0.83	0/2477
1	B	0.60	0/1817	0.86	0/2458
1	C	0.57	0/1823	0.83	0/2466
1	D	0.57	0/1825	0.86	0/2469
All	All	0.58	0/7295	0.85	0/9870

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1790	0	1748	15	0
1	B	1780	0	1749	17	0
1	C	1786	0	1754	17	0
1	D	1788	0	1752	16	0
2	A	109	0	0	1	0
2	B	104	0	0	2	0
2	C	80	0	0	1	0
2	D	92	0	0	2	0
All	All	7529	0	7003	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:68:GLN:HE21	1:D:72:VAL:HG23	1.43	0.83
1:D:66:HIS:CE1	1:D:70:GLN:NE2	2.49	0.81
1:D:88:GLU:OE2	1:D:180:LEU:HD22	1.83	0.78
1:A:89:GLU:HG2	1:A:200:GLN:HE22	1.52	0.74
1:B:177:PRO:O	1:B:178:GLU:HB2	1.97	0.64
1:B:145:LEU:HD12	1:B:268:CYS:HB3	1.81	0.62
1:D:66:HIS:CE1	1:D:70:GLN:HE22	2.17	0.62
1:C:68:GLN:CG	1:C:228:ARG:HH21	2.17	0.58
1:A:282:SER:O	1:A:285:SER:HB3	2.03	0.58
1:A:145:LEU:HD12	1:A:268:CYS:HB3	1.86	0.58
1:C:188:THR:CG2	1:C:190:ASN:O	2.53	0.57
1:A:77:ARG:HB3	1:A:77:ARG:HH11	1.71	0.55
1:B:73:LYS:O	1:B:77:ARG:HG3	2.08	0.54
1:C:90:LYS:NZ	2:C:303:HOH:O	2.40	0.54
1:C:145:LEU:HD12	1:C:268:CYS:HB3	1.90	0.54
1:C:188:THR:HG22	1:C:190:ASN:O	2.08	0.54
1:B:95:LEU:HD21	1:B:181:ILE:HD11	1.89	0.54
1:C:95:LEU:HD21	1:C:181:ILE:HD11	1.89	0.53
1:A:95:LEU:HD21	1:A:181:ILE:HD11	1.90	0.53
1:D:95:LEU:HD21	1:D:181:ILE:HD11	1.93	0.51
1:C:68:GLN:HG3	1:C:228:ARG:HH21	1.76	0.50
1:D:145:LEU:HD12	1:D:268:CYS:HB3	1.93	0.50
1:A:71:GLU:OE1	1:A:228:ARG:CZ	2.59	0.50
1:A:177:PRO:O	1:A:178:GLU:HB2	2.11	0.50
1:C:85:LEU:HB2	1:C:90:LYS:HG3	1.94	0.49
1:B:70:GLN:HE22	1:D:114:ASP:C	2.21	0.49
1:A:77:ARG:HB3	1:A:77:ARG:NH1	2.29	0.47
1:D:66:HIS:CE1	1:D:70:GLN:HE21	2.27	0.47
1:C:65:THR:O	1:C:69:VAL:HG13	2.15	0.47
1:A:73:LYS:HE3	1:A:73:LYS:HB3	1.53	0.47
1:B:200:GLN:NE2	2:B:302:HOH:O	2.33	0.46
1:B:200:GLN:HG2	1:B:201:PHE:CE2	2.50	0.46
1:B:241:GLU:N	1:B:241:GLU:CD	2.74	0.46
1:B:66:HIS:O	1:B:70:GLN:HG2	2.16	0.46
1:B:73:LYS:HG2	1:B:209:LEU:HD11	1.98	0.46
1:D:200:GLN:HG2	1:D:201:PHE:CE2	2.51	0.46
1:A:94:GLU:OE1	1:C:283:ILE:HG13	2.17	0.45
1:C:283:ILE:C	1:C:283:ILE:HD12	2.42	0.45
1:A:65:THR:O	1:A:69:VAL:HG13	2.17	0.45
1:A:91:LYS:HA	1:C:283:ILE:HG23	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:70:GLN:NE2	1:D:113:THR:O	2.50	0.44
1:B:188:THR:HG22	1:B:193:PHE:CD2	2.53	0.44
1:C:189:PHE:HB3	1:C:229:PHE:HB3	1.99	0.44
1:B:282:SER:OG	2:B:301:HOH:O	2.21	0.44
1:C:66:HIS:HA	1:C:112:PRO:HG3	2.01	0.43
1:D:148:ALA:HB1	1:D:165:PHE:HA	2.01	0.43
2:A:313:HOH:O	1:C:282:SER:CB	2.66	0.42
1:B:70:GLN:NE2	1:D:114:ASP:C	2.77	0.42
1:D:73:LYS:NZ	2:D:304:HOH:O	2.48	0.42
1:B:66:HIS:CE1	1:B:70:GLN:NE2	2.87	0.42
1:A:148:ALA:HB1	1:A:165:PHE:HA	2.02	0.42
1:C:145:LEU:CD1	1:C:268:CYS:HB3	2.49	0.42
1:A:145:LEU:CD1	1:A:268:CYS:HB3	2.47	0.42
1:B:145:LEU:CD1	1:B:268:CYS:HB3	2.48	0.42
1:C:188:THR:HG21	1:C:190:ASN:O	2.19	0.42
1:D:200:GLN:NE2	2:D:308:HOH:O	2.53	0.42
1:D:281:LEU:HB2	1:D:284:VAL:HG13	2.02	0.42
1:B:102:LEU:HD12	1:B:169:LEU:HD11	2.02	0.41
1:D:145:LEU:CD1	1:D:268:CYS:HB3	2.50	0.41
1:A:71:GLU:OE1	1:A:228:ARG:NH2	2.54	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	221/231 (96%)	214 (97%)	7 (3%)	0	100	100
1	B	218/231 (94%)	212 (97%)	6 (3%)	0	100	100
1	C	219/231 (95%)	213 (97%)	6 (3%)	0	100	100
1	D	219/231 (95%)	213 (97%)	6 (3%)	0	100	100
All	All	877/924 (95%)	852 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/207 (95%)	191 (97%)	6 (3%)	36	24
1	B	197/207 (95%)	196 (100%)	1 (0%)	86	86
1	C	198/207 (96%)	195 (98%)	3 (2%)	60	53
1	D	197/207 (95%)	193 (98%)	4 (2%)	50	40
All	All	789/828 (95%)	775 (98%)	14 (2%)	58	45

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	ARG
1	A	84	HIS
1	A	88	GLU
1	A	188	THR
1	A	256[A]	GLU
1	A	256[B]	GLU
1	B	218	GLN
1	C	178	GLU
1	C	270	HIS
1	C	282	SER
1	D	77	ARG
1	D	84	HIS
1	D	88	GLU
1	D	273	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	200	GLN
1	B	66	HIS
1	B	70	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	182	HIS
1	C	66	HIS
1	D	66	HIS
1	D	70	GLN
1	D	191	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	222/231 (96%)	-0.10	5 (2%) 61 59	20, 33, 68, 87	1 (0%)
1	B	220/231 (95%)	-0.12	4 (1%) 67 66	20, 33, 67, 82	0
1	C	221/231 (95%)	-0.03	2 (0%) 81 80	22, 36, 68, 90	0
1	D	221/231 (95%)	0.03	5 (2%) 61 59	22, 36, 74, 94	0
All	All	884/924 (95%)	-0.06	16 (1%) 67 66	20, 35, 71, 94	1 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	190	ASN	3.7
1	D	190	ASN	3.4
1	D	284	VAL	3.2
1	D	64	SER	3.1
1	B	252	GLY	3.1
1	C	190	ASN	2.7
1	A	249	VAL	2.5
1	A	252	GLY	2.5
1	B	283	ILE	2.5
1	D	77	ARG	2.4
1	B	113	THR	2.3
1	A	64	SER	2.2
1	B	67	SER	2.1
1	D	283	ILE	2.1
1	A	68	GLN	2.1
1	C	78	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.