



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 22, 2026 – 02:20 AM JST

PDB ID : 9WBE / pdb_00009wbe
Title : Crystal structure of the CHS-CHIL complex
Authors : Li, J.X.; zhang, P.
Deposited on : 2025-08-13
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1028 (1.92-1.92)
Clashscore	180529	1100 (1.92-1.92)
Ramachandran outliers	177936	1087 (1.92-1.92)
Sidechain outliers	177891	1087 (1.92-1.92)
RSRZ outliers	164620	1028 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	89% 9% .
1	B	395	91% 8% .
2	C	209	82% 16% .
2	D	209	79% 19% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 16123 atoms, of which 6046 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chalcone synthase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	389	Total	C	H	N	O	S	0	3	0
			6034	1901	3025	520	570	18			
1	B	391	Total	C	H	N	O	S	0	3	0
			6039	1906	3021	522	572	18			

- Molecule 2 is a protein called Probable chalcone-flavanone isomerase 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	206	Total	C	N	O	S	0	0	0
			1621	1034	260	321	6			
2	D	206	Total	C	N	O	S	0	0	0
			1621	1034	260	321	6			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	310	Total	O	0	0
			310	310		
3	B	294	Total	O	0	0
			294	294		
3	C	111	Total	O	0	0
			111	111		
3	D	93	Total	O	0	0
			93	93		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Chalcone synthase




- Molecule 1: Chalcone synthase



- Molecule 2: Probable chalcone–flavanone isomerase 3



- Molecule 2: Probable chalcone–flavanone isomerase 3





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.46Å 114.91Å 146.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.99 – 1.91 34.99 – 1.91	Depositor EDS
% Data completeness (in resolution range)	100.0 (34.99-1.91) 100.0 (34.99-1.91)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 1.91Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.170 , 0.204 0.170 , 0.204	Depositor DCC
R_{free} test set	4809 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	26.1	Xtriage
Anisotropy	0.166	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16123	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	1/3060 (0.0%)	0.71	2/4139 (0.0%)
1	B	0.53	3/3069 (0.1%)	0.69	0/4151
2	C	0.56	0/1650	0.79	2/2227 (0.1%)
2	D	0.49	0/1650	0.69	2/2227 (0.1%)
All	All	0.52	4/9429 (0.0%)	0.71	6/12744 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	107	VAL	CA-CB	8.44	1.58	1.54
1	B	277	VAL	CA-CB	7.88	1.58	1.54
1	A	96	LEU	C-O	-6.79	1.16	1.24
1	B	96	LEU	C-O	-5.93	1.17	1.24

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	27	GLY	N-CA-C	-9.51	96.31	110.38
2	D	14	PRO	O-C-N	-6.93	118.12	121.31
2	D	20	SER	N-CA-C	-6.72	105.23	113.50
1	A	95	SER	CB-CA-C	5.75	116.57	109.16
2	C	20	SER	N-CA-C	-5.41	106.66	113.20
1	A	96	LEU	CA-C-O	-5.06	115.25	120.92

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3009	3025	3026	23	1
1	B	3018	3021	3034	25	0
2	C	1621	0	1607	25	0
2	D	1621	0	1607	37	1
3	A	310	0	0	8	1
3	B	294	0	0	11	1
3	C	111	0	0	3	0
3	D	93	0	0	2	0
All	All	10077	6046	9274	106	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:49:LEU:HD22	2:C:84:MET:HE2	1.40	1.04
2:D:49:LEU:HD22	2:D:84:MET:HE2	1.40	1.02
1:A:75:MET:SD	3:A:658:HOH:O	2.19	0.97
2:D:49:LEU:HD21	2:D:86:LYS:HG2	1.45	0.95
2:D:167:GLU:HG2	2:D:169:LYS:HE3	1.50	0.93
1:A:169:CSD:OD1	3:A:401:HOH:O	1.87	0.89
2:C:60:ASN:HB3	3:C:303:HOH:O	1.75	0.87
2:C:49:LEU:HD22	2:C:84:MET:CE	2.09	0.82
2:D:49:LEU:HD22	2:D:84:MET:CE	2.13	0.79
2:C:65:THR:HG22	2:C:68:GLU:OE1	1.84	0.78
1:A:287:LYS:NZ	3:A:402:HOH:O	2.05	0.76
1:B:334:ARG:NE	3:B:401:HOH:O	1.90	0.74
2:C:65:THR:HG23	2:C:68:GLU:H	1.53	0.73
2:D:49:LEU:CD2	2:D:86:LYS:HG2	2.17	0.72
1:B:290:ASP:OD2	3:B:402:HOH:O	2.08	0.71
1:A:126:LYS:NZ	3:A:406:HOH:O	2.23	0.71
1:A:255:ASP:OD2	3:A:403:HOH:O	2.10	0.70
1:B:64:MET:HE2	2:C:38:LEU:HD13	1.72	0.69
2:C:86:LYS:NZ	2:C:182:GLN:OE1	2.25	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:113:GLU:OE1	3:B:404:HOH:O	2.12	0.67
1:B:19:ASP:OD2	3:B:405:HOH:O	2.14	0.66
2:C:111:ASP:O	2:C:115:GLU:HG3	1.95	0.65
2:C:128:GLU:OE2	2:C:128:GLU:HA	1.97	0.65
2:D:167:GLU:CG	2:D:169:LYS:HE3	2.26	0.64
2:D:125:THR:HG22	2:D:129:LYS:NZ	2.13	0.63
2:D:206:ALA:O	2:D:209:THR:HG22	1.98	0.63
1:B:187:ARG:NH2	3:B:403:HOH:O	2.12	0.61
2:D:126:GLU:HA	2:D:129:LYS:HE2	1.81	0.61
2:D:66:GLY:HA3	2:D:202:ASP:OD1	2.01	0.61
2:D:183:ARG:O	2:D:187:SER:HB2	2.01	0.60
1:B:57:LEU:HD11	1:B:214:LEU:HD11	1.83	0.60
2:D:112:ARG:HB2	2:D:180:MET:HE2	1.84	0.59
1:A:276:ASP:N	2:D:33:ILE:HD11	2.18	0.59
1:B:187:ARG:NE	3:B:403:HOH:O	2.30	0.59
1:B:64:MET:CE	2:C:38:LEU:HD13	2.32	0.58
2:D:125:THR:HG22	2:D:129:LYS:HZ1	1.67	0.57
2:D:128:GLU:HG3	3:D:355:HOH:O	2.04	0.56
2:D:137:LYS:HE3	2:D:168:GLU:OE1	2.05	0.56
2:D:167:GLU:HG2	2:D:169:LYS:CE	2.31	0.56
2:C:105:LEU:C	2:C:105:LEU:HD23	2.32	0.55
2:D:149:PHE:CE1	2:D:182:GLN:HG3	2.43	0.54
2:C:60:ASN:ND2	3:C:303:HOH:O	2.40	0.54
1:A:127:SER:O	1:A:156:ARG:HD2	2.09	0.52
1:A:162:LEU:HD22	1:B:178:ILE:HD11	1.91	0.51
2:D:140:LYS:NZ	2:D:163:GLU:O	2.44	0.51
2:C:49:LEU:HD21	2:C:86:LYS:HG2	1.91	0.51
1:A:167:GLN:HB2	3:A:618:HOH:O	2.10	0.50
2:C:30:ILE:HD12	2:C:32:ASP:HB3	1.93	0.50
2:D:135:GLN:HB3	3:D:369:HOH:O	2.12	0.50
2:D:126:GLU:HA	2:D:129:LYS:CE	2.41	0.50
1:A:57:LEU:HD11	1:A:211:LEU:CD2	2.42	0.49
2:C:60:ASN:HB2	3:C:334:HOH:O	2.13	0.49
2:D:155:ILE:CD1	2:D:173:GLU:HA	2.43	0.49
1:B:57:LEU:HD11	1:B:214:LEU:CD1	2.43	0.48
2:D:155:ILE:HD13	2:D:173:GLU:HA	1.95	0.48
2:D:149:PHE:CD1	2:D:182:GLN:HG3	2.48	0.48
1:A:113:GLU:OE1	3:A:404:HOH:O	2.20	0.47
1:A:246:VAL:HG21	1:A:373:TRP:HZ3	1.79	0.47
2:D:114:ALA:C	2:D:116:GLU:H	2.22	0.47
2:C:144:VAL:HB	2:C:161:GLU:HG2	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:66:ASP:O	3:B:406:HOH:O	2.20	0.47
1:B:60:LYS:O	1:B:64:MET:HG3	2.15	0.47
2:D:112:ARG:HD3	2:D:180:MET:CE	2.44	0.47
1:B:334:ARG:NH2	3:B:401:HOH:O	2.48	0.47
1:A:291:GLU:OE2	3:A:405:HOH:O	2.21	0.46
1:A:137:THR:O	1:A:137:THR:HG22	2.16	0.45
1:A:193:VAL:O	1:A:226:ALA:HA	2.16	0.45
1:B:376:LEU:C	1:B:376:LEU:HD23	2.41	0.45
2:D:112:ARG:HD3	2:D:180:MET:HE1	1.99	0.45
1:B:334:ARG:CZ	3:B:401:HOH:O	2.51	0.45
1:A:199:THR:HG23	1:A:222:ASP:OD2	2.16	0.45
1:A:64:MET:HE1	1:A:214:LEU:C	2.43	0.44
1:A:64:MET:HE1	1:A:215:VAL:N	2.33	0.44
1:B:109:LYS:HB2	1:B:109:LYS:NZ	2.32	0.44
2:D:158:ILE:HD11	2:D:181:MET:HE1	2.00	0.43
2:D:180:MET:HE3	2:D:180:MET:HA	2.00	0.43
1:B:121:GLU:CD	1:B:356:ARG:HH22	2.26	0.43
2:D:95:GLU:HG3	2:D:140:LYS:HA	1.99	0.43
2:D:145:ILE:HD12	2:D:145:ILE:N	2.33	0.43
1:B:272:HIS:HB3	1:B:274:LEU:HD11	2.00	0.43
1:B:294:LYS:O	1:B:294:LYS:HD3	2.19	0.43
2:C:65:THR:CG2	2:C:67:LYS:HG3	2.49	0.43
2:C:116:GLU:OE2	2:C:118:LYS:HE2	2.18	0.43
1:A:197:GLU:HG3	1:A:344:SER:HB3	2.01	0.43
2:D:161:GLU:HG2	2:D:167:GLU:CD	2.44	0.43
1:B:47:ARG:NH2	3:B:409:HOH:O	2.39	0.42
2:C:110:ARG:HG3	2:C:119:TYR:CZ	2.55	0.42
2:D:137:LYS:HG2	2:D:160:PHE:CE2	2.55	0.42
2:C:76:PHE:CZ	2:C:197:ILE:HG23	2.55	0.42
1:B:187:ARG:CZ	3:B:403:HOH:O	2.55	0.42
1:B:207:SER:HB3	1:B:210:HIS:CE1	2.55	0.42
2:C:25:LEU:HD13	2:C:48:TYR:CZ	2.55	0.42
1:A:292:ALA:HB1	1:A:389:VAL:CG2	2.49	0.42
2:C:126:GLU:OE1	2:C:129:LYS:HE2	2.20	0.41
2:D:33:ILE:CG2	2:D:35:ILE:HD11	2.51	0.41
1:B:382:GLY:N	1:B:383:LEU:HA	2.36	0.41
2:C:54:VAL:CG1	2:C:204:ILE:HG23	2.50	0.41
1:B:193:VAL:O	1:B:226:ALA:HA	2.21	0.41
1:A:258:GLY:O	1:A:274:LEU:HB2	2.21	0.41
1:A:83:LYS:HB3	1:A:83:LYS:HE2	1.86	0.40
1:A:307:ILE:O	1:A:376:LEU:HA	2.22	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:26:LEU:HD23	2:D:26:LEU:HA	1.86	0.40
2:D:155:ILE:HD13	2:D:155:ILE:HA	1.87	0.40
2:C:33:ILE:HG21	2:C:35:ILE:HD11	2.04	0.40
2:D:35:ILE:O	2:D:38:LEU:HB3	2.20	0.40
2:C:65:THR:HG23	2:C:67:LYS:HG3	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:120:LYS:HZ1	2:D:173:GLU:OE1[1_655]	1.49	0.11
3:A:666:HOH:O	3:B:563:HOH:O[3_555]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	389/395 (98%)	379 (97%)	10 (3%)	0	100	100
1	B	391/395 (99%)	384 (98%)	7 (2%)	0	100	100
2	C	204/209 (98%)	195 (96%)	9 (4%)	0	100	100
2	D	204/209 (98%)	198 (97%)	6 (3%)	0	100	100
All	All	1188/1208 (98%)	1156 (97%)	32 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	327/327 (100%)	327 (100%)	0	100	100
1	B	327/327 (100%)	326 (100%)	1 (0%)	91	90
2	C	179/181 (99%)	177 (99%)	2 (1%)	70	63
2	D	179/181 (99%)	177 (99%)	2 (1%)	70	63
All	All	1012/1016 (100%)	1007 (100%)	5 (0%)	86	85

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	89	CYS
2	C	152	LYS
2	C	161	GLU
2	D	18	ILE
2	D	187	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	167	GLN
1	B	14	GLN
1	B	284	ASN
1	B	335	HIS
2	C	57	HIS
2	C	104	GLN
2	D	16	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSD	B	169	1	3,7,8	0.66	0	1,8,10	3.07	1 (100%)
1	CSD	A	169	1	3,7,8	0.80	0	1,8,10	1.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	B	169	1	-	0/2/6/8	-
1	CSD	A	169	1	-	0/2/6/8	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	169	CSD	OD1-SG-CB	-3.07	99.69	105.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	169	CSD	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/395 (98%)	-0.56	3 (0%) 82 87	12, 27, 50, 85	3 (0%)
1	B	390/395 (98%)	-0.42	5 (1%) 74 80	12, 28, 58, 75	3 (0%)
2	C	206/209 (98%)	0.23	2 (0%) 79 84	22, 38, 64, 89	0
2	D	206/209 (98%)	0.33	5 (2%) 59 65	24, 40, 65, 89	0
All	All	1190/1208 (98%)	-0.23	15 (1%) 74 80	12, 31, 59, 89	6 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	27	GLY	4.2
1	B	91	TYR	3.9
2	D	209	THR	3.8
2	D	63	GLY	3.1
1	A	7	SER	2.9
1	B	6	ALA	2.9
1	B	5	GLY	2.7
2	D	188	GLY	2.6
1	B	48	ILE	2.2
2	C	209	THR	2.2
2	D	37	PHE	2.2
2	D	155	ILE	2.1
1	A	156	ARG	2.1
1	A	11	GLU	2.1
1	B	92	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	CSD	B	169	8/9	0.96	0.07	24,25,33,34	0
1	CSD	A	169	8/9	0.97	0.06	19,24,29,36	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.