



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 25, 2026 – 12:37 PM JST

PDB ID : 9WEL / pdb_00009wel
Title : Plasmodium vivax aspartyl-tRNA synthetase in complex with Asp-AMS, Mg ion, Ethylene glycols and Unit cell contraction due to data collected outside the cryo loop
Authors : Manickam, Y.; Sharma, V.K.; Bagale, S.; Pradeepkumar, P.I.; Sharma, A.
Deposited on : 2025-08-19
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	536	5% 82% 11% • 6%
1	B	536	5% 80% 13% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	PEG	A	704	-	-	-	X

2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 8617 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called aspartate-tRNA ligase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	503	Total	C	N	O	S	0	5	0
			4084	2619	679	758	28			
1	B	501	Total	C	N	O	S	0	2	0
			3990	2561	663	739	27			

- Molecule 2 is 5'-O-(L-alpha-aspartylsulfamoyl)adenosine (CCD ID: DSZ) (formula: C₁₄H₁₉N₇O₉S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		
4	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is BETA-MERCAPTOETHANOL (CCD ID: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	O	S	0	0
			4	2	1	1		
6	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Cl	0	0
			2	2		
7	B	5	Total	Cl	0	0
			5	5		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	191	Total	O	0	27
			218	218		
8	B	175	Total	O	0	9
			184	184		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: aspartate-tRNA ligase



• Molecule 1: aspartate-tRNA ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	136.48Å 136.48Å 273.35Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	89.40 – 2.20 89.40 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.3 (89.40-2.20) 99.3 (89.40-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 2.20Å)	Xtriage
Refinement program	PHENIX (1.15rc1_3423: ???)	Depositor
R, R_{free}	0.178 , 0.214 0.190 , 0.218	Depositor DCC
R_{free} test set	3849 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	55.4	Xtriage
Anisotropy	0.136	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 78.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8617	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, CL, BME, PEG, DSZ, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.37	0/4186	0.66	6/5646 (0.1%)
1	B	0.39	0/4086	0.57	1/5524 (0.0%)
All	All	0.38	0/8272	0.62	7/11170 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	228[A]	MET	CA-C-O	13.25	134.47	120.42
1	A	228[B]	MET	CA-C-O	13.25	134.47	120.42
1	A	228[A]	MET	N-CA-C	9.50	121.72	111.36
1	A	228[B]	MET	N-CA-C	9.50	121.72	111.36
1	A	289	THR	N-CA-C	5.59	115.50	108.45
1	B	255	SER	N-CA-C	5.05	117.90	111.69
1	A	125	LYS	N-CA-C	-5.02	101.53	109.96

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4084	0	3952	48	0
1	B	3990	0	3803	47	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	31	0	18	0	0
2	B	31	0	18	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	10	0	14	0	0
4	B	10	0	14	0	0
5	A	28	0	40	4	0
5	B	14	0	20	3	0
6	A	4	0	6	0	0
6	B	4	0	6	0	0
7	A	2	0	0	0	0
7	B	5	0	0	0	0
8	A	218	0	0	8	0
8	B	184	0	0	2	0
All	All	8617	0	7891	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:404:ARG:NH2	1:A:605:ARG:HH12	1.54	1.05
1:B:219:LYS:HA	1:B:223:ASN:ND2	1.78	0.98
1:A:228[B]:MET:CE	8:A:844:HOH:O	2.30	0.78
1:A:228[B]:MET:HE2	8:A:844:HOH:O	1.85	0.76
1:B:219:LYS:HA	1:B:223:ASN:HD22	1.50	0.76
1:A:404:ARG:NH2	1:A:605:ARG:NH1	2.37	0.70
1:A:463:GLN:HA	5:A:707:PEG:H12	1.73	0.70
5:A:707:PEG:H21	8:A:971[B]:HOH:O	1.97	0.64
1:A:358:PHE:HB3	1:B:360:ILE:HD11	1.80	0.64
1:B:219:LYS:HA	1:B:223:ASN:HD21	1.59	0.64
1:A:404:ARG:HH21	1:A:605:ARG:HH12	1.44	0.63
1:B:297:VAL:HG12	1:B:302:ARG:HG3	1.81	0.63
1:B:525:PRO:HG2	1:B:528:LEU:HD12	1.82	0.60
1:A:228[B]:MET:HE1	8:A:844:HOH:O	1.97	0.60
1:A:527:ALA:HA	5:A:705:PEG:H41	1.85	0.58
1:B:477:ILE:HG21	1:B:493:ILE:HG23	1.86	0.58
1:A:508:LYS:NZ	8:A:896[B]:HOH:O	2.36	0.57
1:A:197:LYS:HD2	1:A:202:PHE:CE1	2.39	0.57
1:B:496:TYR:HA	5:B:705:PEG:H42	1.86	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:487:HIS:O	1:A:488:LEU:HG	2.05	0.56
1:A:591:TYR:HA	1:B:273:THR:HG23	1.88	0.56
1:B:104:GLU:O	1:B:108:VAL:HG23	2.05	0.56
1:B:277:LEU:H	1:B:277:LEU:HD12	1.71	0.55
1:B:132:ASN:O	1:B:134:LYS:N	2.33	0.55
1:A:584:PHE:HD2	1:A:584:PHE:H	1.53	0.55
1:A:289:THR:HG23	1:A:291:GLU:H	1.74	0.53
1:A:348:GLY:O	1:A:368:PHE:HD2	1.93	0.52
1:B:141:LEU:HB2	1:B:207:HIS:HB3	1.92	0.51
1:A:116:ILE:HD11	1:A:130:LYS:C	2.35	0.51
1:A:213:GLN:NE2	1:A:261:GLU:OE1	2.41	0.51
1:A:117:ASN:HD21	1:A:131:GLU:HA	1.77	0.50
1:B:282:LYS:O	1:B:286:MET:HG3	2.12	0.50
1:B:136:ILE:HG23	1:B:186:TRP:CD1	2.46	0.49
1:B:192:HIS:HB2	1:B:206:ARG:HB2	1.94	0.49
1:B:522:ILE:HG22	1:B:546:TYR:HB3	1.95	0.49
1:B:133:GLU:N	1:B:133:GLU:OE1	2.46	0.49
1:A:376:TYR:HA	1:A:379:MET:HE3	1.94	0.49
1:B:487:HIS:O	1:B:488:LEU:HD23	2.14	0.48
1:B:291:GLU:O	1:B:294:SER:CB	2.62	0.48
1:B:349:GLU:HG2	1:B:581:LYS:HB3	1.95	0.48
1:B:460:GLU:HG3	8:B:825:HOH:O	2.12	0.48
1:A:548:PHE:HB2	1:A:556:LEU:HB3	1.97	0.47
1:B:217:ASP:OD2	1:B:219:LYS:HG2	2.15	0.47
1:B:291:GLU:CB	1:B:294:SER:OG	2.63	0.47
1:B:465:LEU:HG	1:B:516:THR:HB	1.96	0.47
1:B:554:GLU:O	1:B:605:ARG:NH2	2.47	0.47
1:A:477:ILE:O	1:A:481:ILE:HG13	2.15	0.47
1:B:548:PHE:HB2	1:B:556:LEU:HB3	1.97	0.46
1:A:197:LYS:HD2	1:A:202:PHE:HE1	1.80	0.46
1:A:276:GLU:OE2	1:A:276:GLU:HA	2.16	0.46
1:A:192:HIS:HB2	1:A:206:ARG:HB2	1.98	0.46
1:A:302:ARG:HB3	1:A:309:ASP:CG	2.41	0.46
1:A:99:GLU:OE2	1:A:99:GLU:HA	2.16	0.45
1:A:481:ILE:HA	1:A:486:LEU:O	2.15	0.45
1:A:348:GLY:C	1:A:350:SER:H	2.25	0.45
5:A:706:PEG:H11	8:A:823:HOH:O	2.15	0.45
5:B:705:PEG:H12	5:B:705:PEG:H31	1.76	0.45
1:B:115:ASP:O	1:B:119:GLU:HG3	2.16	0.45
1:B:496:TYR:CE2	1:B:498:MET:HA	2.52	0.45
1:A:584:PHE:N	1:A:584:PHE:CD2	2.85	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:LEU:O	1:A:211:SER:HA	2.17	0.44
1:A:121:TYR:HA	1:A:267:ILE:O	2.18	0.44
1:B:513:SER:OG	1:B:514:HIS:ND1	2.44	0.44
1:B:216:LEU:HD11	1:B:221:ASN:ND2	2.33	0.44
1:A:218:ILE:HG12	1:A:223:ASN:HA	2.00	0.44
1:B:326:CYS:O	1:B:330:LYS:HE3	2.17	0.44
1:B:217:ASP:CG	1:B:219:LYS:HG2	2.42	0.43
1:B:281:LEU:HD23	1:B:281:LEU:HA	1.90	0.43
1:A:520:ILE:HG12	1:A:548:PHE:CD1	2.54	0.43
1:B:450:ILE:O	1:B:454:LYS:HG2	2.18	0.43
1:B:120:SER:HA	1:B:266:LYS:CD	2.48	0.43
1:B:583:ASN:OD1	1:B:583:ASN:N	2.51	0.43
1:A:218:ILE:HD12	1:A:228[B]:MET:HG3	2.01	0.43
1:A:194:ILE:HD13	1:A:194:ILE:HG21	1.75	0.42
1:A:470:ILE:HD11	8:A:972:HOH:O	2.19	0.42
1:B:291:GLU:O	1:B:294:SER:OG	2.37	0.42
1:A:194:ILE:HG12	1:A:203:ILE:HG12	2.01	0.42
1:A:348:GLY:HA3	1:B:630:ILE:HD11	2.02	0.42
1:A:523:ASN:HA	1:A:543:SER:O	2.19	0.42
1:A:307[B]:CYS:HB3	8:A:929:HOH:O	2.19	0.42
1:A:386:ASP:OD2	5:B:706:PEG:H12	2.20	0.42
1:B:504:LYS:HB2	1:B:504:LYS:HE3	1.70	0.42
1:B:523:ASN:HA	1:B:543:SER:O	2.20	0.42
1:A:140:ASN:OD1	1:A:143:GLU:HG3	2.21	0.41
1:B:214:CYS:HA	1:B:262:ALA:O	2.20	0.41
1:A:440:LEU:HD23	1:A:440:LEU:HA	1.93	0.41
1:A:211:SER:OG	1:A:256:THR:HA	2.20	0.41
1:A:535:TYR:HA	1:A:543:SER:HA	2.03	0.41
1:B:265:ARG:C	1:B:266:LYS:HG2	2.46	0.41
1:A:235:PRO:HG2	1:A:273:THR:HB	2.03	0.41
1:B:398:GLU:HB3	8:B:832:HOH:O	2.20	0.41
1:B:404:ARG:NH1	1:B:605:ARG:NH2	2.69	0.41
1:B:374:GLN:HG2	1:B:532:TYR:CE1	2.57	0.40
1:A:465:LEU:HG	1:A:516:THR:HB	2.02	0.40
1:B:359:GLN:HG2	1:B:368:PHE:CE2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	500/536 (93%)	484 (97%)	16 (3%)	0	100	100
1	B	499/536 (93%)	482 (97%)	16 (3%)	1 (0%)	43	51
All	All	999/1072 (93%)	966 (97%)	32 (3%)	1 (0%)	48	57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	129	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	441/485 (91%)	440 (100%)	1 (0%)	87	94
1	B	421/485 (87%)	421 (100%)	0	100	100
All	All	862/970 (89%)	861 (100%)	1 (0%)	88	95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	218	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	ASN
1	A	299	GLN
1	A	580	ASN
1	B	207	HIS
1	B	220	HIS
1	B	221	ASN
1	B	223	ASN
1	B	456	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 9 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PEG	A	706	-	6,6,6	0.36	0	5,5,5	0.60	0
5	PEG	A	707	-	6,6,6	0.15	0	5,5,5	0.19	0
2	DSZ	B	702	-	32,33,33	0.18	0	45,49,49	0.31	0
2	DSZ	A	701	-	32,33,33	0.17	0	45,49,49	0.30	0
6	BME	B	707	-	3,3,3	0.27	0	1,2,2	0.14	0
4	PGE	B	704	-	9,9,9	0.33	0	8,8,8	0.27	0
5	PEG	A	704	-	6,6,6	0.43	0	5,5,5	0.31	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	PEG	A	705	-	6,6,6	0.43	0	5,5,5	0.31	0
5	PEG	B	705	-	6,6,6	0.15	0	5,5,5	0.06	0
6	BME	A	708	-	3,3,3	0.35	0	1,2,2	0.22	0
5	PEG	B	706	-	6,6,6	0.43	0	5,5,5	0.31	0
4	PGE	A	703	-	9,9,9	0.34	0	8,8,8	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PEG	A	706	-	-	3/4/4/4	-
5	PEG	A	707	-	-	2/4/4/4	-
2	DSZ	B	702	-	-	6/22/39/39	0/3/3/3
2	DSZ	A	701	-	-	4/22/39/39	0/3/3/3
6	BME	B	707	-	-	0/1/1/1	-
4	PGE	B	704	-	-	3/7/7/7	-
5	PEG	A	704	-	-	0/4/4/4	-
5	PEG	A	705	-	-	0/4/4/4	-
5	PEG	B	705	-	-	1/4/4/4	-
6	BME	A	708	-	-	0/1/1/1	-
5	PEG	B	706	-	-	0/4/4/4	-
4	PGE	A	703	-	-	2/7/7/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	703	PGE	O2-C3-C4-O3
5	A	706	PEG	O1-C1-C2-O2
5	B	705	PEG	C1-C2-O2-C3
5	A	706	PEG	O2-C3-C4-O4
2	A	701	DSZ	O-C-CA-N
2	A	701	DSZ	NAT-C-CA-N
2	B	702	DSZ	O-C-CA-N
2	B	702	DSZ	NAT-C-CA-N

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	701	DSZ	O-C-CA-CB
4	B	704	PGE	O1-C1-C2-O2
5	A	706	PEG	C1-C2-O2-C3
4	B	704	PGE	C4-C3-O2-C2
4	A	703	PGE	C1-C2-O2-C3
2	A	701	DSZ	NAT-C-CA-CB
2	B	702	DSZ	O-C-CA-CB
2	B	702	DSZ	NAT-C-CA-CB
5	A	707	PEG	O1-C1-C2-O2
2	B	702	DSZ	CA-CB-CG-OD2
5	A	707	PEG	C4-C3-O2-C2
2	B	702	DSZ	CA-CB-CG-OD1
4	B	704	PGE	O2-C3-C4-O3

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	706	PEG	1	0
5	A	707	PEG	2	0
5	A	705	PEG	1	0
5	B	705	PEG	2	0
5	B	706	PEG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	503/536 (93%)	0.14	26 (5%)	33 29	34, 60, 107, 145	5 (0%)
1	B	501/536 (93%)	0.29	28 (5%)	30 27	31, 64, 118, 164	3 (0%)
All	All	1004/1072 (93%)	0.21	54 (5%)	31 28	31, 62, 113, 164	8 (0%)

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	350	SER	9.1
1	B	253	ILE	6.1
1	B	116	ILE	5.8
1	B	180	LEU	5.6
1	A	352	GLU	5.4
1	B	179	HIS	4.7
1	B	117	ASN	4.5
1	A	98	ARG	4.3
1	B	295	ILE	4.3
1	B	130	LYS	4.2
1	B	353	GLY	4.1
1	B	115	ASP	4.0
1	A	253	ILE	3.9
1	A	488	LEU	3.9
1	B	492	GLU	3.9
1	B	132	ASN	3.8
1	A	348	GLY	3.8
1	A	487	HIS	3.7
1	B	290	ASN	3.6
1	B	135	GLU	3.4
1	A	295	ILE	3.1
1	B	149	MET	3.1
1	A	251	VAL	3.1
1	B	102	LYS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	254	ASP	3.0
1	A	584	PHE	3.0
1	A	149	MET	3.0
1	A	250	GLU	2.9
1	A	179	HIS	2.9
1	B	255	SER	2.7
1	A	494	LEU	2.7
1	A	252	PRO	2.7
1	A	178	ALA	2.6
1	A	290	ASN	2.6
1	B	291	GLU	2.6
1	B	196	SER	2.6
1	A	486	LEU	2.5
1	A	218	ILE	2.5
1	B	105	ALA	2.4
1	B	491	GLU	2.4
1	B	218	ILE	2.4
1	A	296	LYS	2.4
1	A	222	ASP	2.4
1	A	398[A]	GLU	2.4
1	B	251	VAL	2.3
1	B	195	ARG	2.2
1	B	354	GLY	2.2
1	B	263	HIS	2.2
1	A	219	LYS	2.1
1	A	403	TYR	2.1
1	B	247	SER	2.1
1	A	355	ALA	2.1
1	B	106	THR	2.0
1	A	130	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	PEG	B	705	7/7	0.76	0.21	107,114,116,117	0
5	PEG	A	705	7/7	0.77	0.23	93,103,105,107	7
5	PEG	A	704	7/7	0.79	0.48	85,97,102,102	7
4	PGE	B	704	10/10	0.84	0.23	108,110,113,113	0
5	PEG	A	706	7/7	0.84	0.15	95,102,108,109	0
4	PGE	A	703	10/10	0.84	0.16	74,93,102,106	0
6	BME	A	708	4/4	0.84	0.26	66,71,73,77	4
5	PEG	B	706	7/7	0.85	0.25	99,101,105,106	0
5	PEG	A	707	7/7	0.85	0.23	76,79,83,84	7
7	CL	A	709	1/1	0.88	0.09	109,109,109,109	0
6	BME	B	707	4/4	0.90	0.20	64,67,71,74	4
7	CL	B	711	1/1	0.90	0.15	95,95,95,95	1
7	CL	B	710	1/1	0.91	0.19	93,93,93,93	1
7	CL	B	701	1/1	0.91	0.20	110,110,110,110	0
7	CL	B	709	1/1	0.92	0.20	103,103,103,103	0
3	MG	A	702	1/1	0.93	0.07	83,83,83,83	1
7	CL	B	708	1/1	0.95	0.14	107,107,107,107	0
7	CL	A	710	1/1	0.96	0.15	102,102,102,102	0
3	MG	B	703	1/1	0.97	0.04	65,65,65,65	1
2	DSZ	B	702	31/31	0.98	0.06	40,45,49,51	0
2	DSZ	A	701	31/31	0.98	0.07	43,48,53,55	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around DSZ B 702:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around DSZ A 701:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.