



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 22, 2025 – 06:27 PM JST

PDB ID : 9X2Q / pdb_00009x2q
Title : GSK3beta complexed with BiS-1
Authors : Haslboeck, S.; Vinogradov, A.A.; Okada, C.; Aikawa, H.; Suga, H.; Sengoku, T.
Deposited on : 2025-10-07
Resolution : 1.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.011 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	8422 (1.70-1.66)
Clashscore	180529	1005 (1.68-1.68)
Ramachandran outliers	177936	9065 (1.70-1.66)
Sidechain outliers	177891	9064 (1.70-1.66)
RSRZ outliers	164620	8421 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	359	
1	C	359	
2	B	17	
2	D	17	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PRO	A	502	-	-	X	-
4	PRO	C	502	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6385 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycogen synthase kinase-3 beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	349	Total	C	N	O	S	0	4	0
			2824	1815	484	511	14			
1	C	344	Total	C	N	O	S	0	7	0
			2805	1802	480	508	15			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	GLY	-	expression tag	UNP P49841
A	26	PRO	-	expression tag	UNP P49841
C	25	GLY	-	expression tag	UNP P49841
C	26	PRO	-	expression tag	UNP P49841


- Molecule 2 is a protein called BiS-1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	10	Total	C	N	O	0	0	0
			120	79	25	16			
2	D	10	Total	C	N	O	0	0	0
			120	79	25	16			

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	C	1	Total	Cl	0	0
			1	1		

- Molecule 4 is PROLINE (CCD ID: PRO) (formula: C₅H₉NO₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			8	5	1	2		
4	C	1	Total	C	N	O	0	0
			8	5	1	2		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	275	Total 275	O 275	0	0
6	B	6	Total 6	O 6	0	0
6	C	205	Total 205	O 205	0	0
6	D	5	Total 5	O 5	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glycogen synthase kinase-3 beta




- Molecule 1: Glycogen synthase kinase-3 beta



- Molecule 2: BiS-1



- Molecule 2: BiS-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α , β , γ	88.74Å 92.17Å 119.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.58 – 1.68 49.58 – 1.68	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.58-1.68) 100.0 (49.58-1.68)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 1.68Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.173 , 0.202 0.188 , 0.215	Depositor DCC
R_{free} test set	1993 reflections (1.78%)	wwPDB-VP
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.123	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6385	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PEG, A1MCG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	0/2895	0.82	0/3937
1	C	0.68	0/2875	0.76	1/3911 (0.0%)
2	B	0.55	0/65	0.43	0/82
2	D	0.58	0/65	0.45	0/82
All	All	0.73	0/5900	0.79	1/8012 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	C	189	LEU	CB-CG-CD1	-6.91	89.97	110.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	92	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2824	0	2848	36	0
1	C	2805	0	2819	13	0
2	B	120	0	57	2	0
2	D	120	0	57	1	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
4	A	8	0	7	5	0
4	C	8	0	7	5	0
5	A	7	0	10	0	0
6	A	275	0	0	9	1
6	B	6	0	0	0	0
6	C	205	0	0	1	1
6	D	5	0	0	0	0
All	All	6385	0	5805	51	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:502:PRO:CB	4:C:502:PRO:CG	1.74	1.46
4:A:502:PRO:CB	4:A:502:PRO:CG	1.96	1.35
1:A:96:ARG:HH22	4:A:502:PRO:HB3	1.28	0.97
1:C:90:ASP:OD1	1:C:92:ARG:HD3	1.78	0.83
1:A:103:LYS:NZ	6:A:601:HOH:O	2.19	0.75
1:A:373:LEU:HD12	1:A:374:ALA:H	1.55	0.72
4:A:502:PRO:HD2	1:C:292:LYS:H	1.57	0.68
1:C:107[A]:CYS:SG	6:C:617:HOH:O	2.54	0.66
1:A:122:LYS:NZ	6:A:605:HOH:O	2.28	0.65
1:A:111:ARG:NH1	6:A:603:HOH:O	2.20	0.63
1:A:278:ARG:HH12	1:A:290:GLU:CD	2.07	0.63
6:A:687:HOH:O	4:C:502:PRO:HB2	2.00	0.62
1:A:374:ALA:HA	1:A:377:LEU:HB2	1.83	0.61
1:A:376:ILE:N	1:A:376:ILE:HD12	2.19	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:122:LYS:HZ1	1:C:297:LYS:HG2	1.71	0.56
1:C:62:ILE:HD13	1:C:72:GLN:HB2	1.90	0.54
1:A:278:ARG:NH1	1:A:290:GLU:OE2	2.41	0.53
1:A:46:GLN:NE2	1:A:111:ARG:HE	2.05	0.52
1:A:89:GLN:OE1	6:A:602:HOH:O	2.19	0.52
1:C:349:LYS:NZ	1:C:355:ASP:OD1	2.39	0.50
1:A:94:LYS:HD3	1:A:99:GLN:NE2	2.26	0.50
1:C:217:ILE:O	1:C:218[B]:CYS:HB2	2.12	0.50
1:A:122:LYS:NZ	1:C:297:LYS:HG2	2.27	0.50
1:A:220:ARG:HG2	1:A:221:TYR:CD1	2.47	0.49
1:A:96:ARG:NH2	4:A:502:PRO:HB3	2.12	0.48
1:A:373:LEU:HD12	1:A:374:ALA:N	2.25	0.48
1:A:292:LYS:H	4:C:502:PRO:HG3	1.77	0.48
1:A:205:LYS:NZ	4:A:502:PRO:HB2	2.29	0.48
1:C:145:HIS:HB2	2:D:13:TYR:HB3	1.96	0.47
6:A:605:HOH:O	1:C:297:LYS:HG2	2.15	0.47
1:A:94:LYS:HD3	1:A:99:GLN:HE21	1.79	0.47
1:C:97:GLU:O	1:C:101[A]:MET:HG2	2.13	0.47
1:A:217:ILE:O	1:A:218[B]:CYS:HB2	2.14	0.47
1:A:279:GLU:OE2	6:A:604:HOH:O	2.20	0.47
1:A:36:LYS:NZ	6:A:616:HOH:O	2.47	0.47
1:A:39:THR:HG22	1:A:55:SER:OG	2.16	0.46
1:A:292:LYS:H	4:C:502:PRO:CG	2.29	0.46
1:C:59:THR:HA	1:C:72:GLN:O	2.16	0.46
1:A:140:TYR:HB2	1:A:184:PRO:HB2	1.98	0.45
1:A:350:LEU:HD12	1:A:354:ARG:HB2	1.98	0.45
1:A:44:PRO:HG3	1:A:52:GLN:NE2	2.32	0.45
1:A:186:ASN:OD1	2:B:7:ARG:NH2	2.48	0.44
1:A:144:ARG:NE	6:A:619:HOH:O	2.49	0.44
1:A:291:PHE:CD1	4:C:502:PRO:HB3	2.53	0.43
2:B:9:A1MCG:CE2	2:B:9:A1MCG:N15	2.81	0.43
1:C:350:LEU:HD12	1:C:354:ARG:HB2	2.01	0.43
1:A:221:TYR:CD1	1:A:258:PRO:HA	2.54	0.43
1:A:218[B]:CYS:HB3	1:A:223:ARG:HG2	2.02	0.41
1:A:373:LEU:O	1:A:374:ALA:CB	2.68	0.41
1:A:370:ASN:C	1:A:370:ASN:HD22	2.28	0.40
1:A:97:GLU:O	1:A:101[A]:MET:HG2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:767:HOH:O	6:C:755:HOH:O[4_444]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	351/359 (98%)	339 (97%)	10 (3%)	2 (1%)	22	9
1	C	347/359 (97%)	339 (98%)	6 (2%)	2 (1%)	22	9
2	B	5/17 (29%)	5 (100%)	0	0	100	100
2	D	5/17 (29%)	5 (100%)	0	0	100	100
All	All	708/752 (94%)	688 (97%)	16 (2%)	4 (1%)	22	9

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	374	ALA
1	C	49	ASP
1	C	221	TYR
1	A	221	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	315/319 (99%)	310 (98%)	5 (2%)	58	41
1	C	314/319 (98%)	311 (99%)	3 (1%)	73	62

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	7/11 (64%)	7 (100%)	0	100	100
2	D	7/11 (64%)	7 (100%)	0	100	100
All	All	643/660 (97%)	635 (99%)	8 (1%)	65	54

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	ASN
1	A	66	SER
1	A	121	GLU
1	A	155	VAL
1	A	373	LEU
1	C	119	SER
1	C	124	ASP
1	C	154	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	46	GLN
1	A	99	GLN
1	A	265	GLN
1	A	299	HIS
1	A	370	ASN
1	C	64	ASN
1	C	89	GLN
1	C	151	GLN
1	C	295	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	A1MCG	B	9	2	18,19,20	1.61	3 (16%)	21,24,26	2.71	5 (23%)
2	A1MCG	D	8	2	18,19,20	1.43	2 (11%)	21,24,26	2.06	7 (33%)
2	A1MCG	D	6	2	18,19,20	1.66	3 (16%)	21,24,26	2.91	6 (28%)
2	A1MCG	B	8	2	18,19,20	1.40	1 (5%)	21,24,26	2.32	10 (47%)
2	A1MCG	B	6	2	18,19,20	1.57	3 (16%)	21,24,26	2.06	5 (23%)
2	A1MCG	D	9	2	18,19,20	1.63	3 (16%)	21,24,26	2.68	6 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1MCG	B	9	2	-	2/9/10/12	0/2/2/2
2	A1MCG	D	8	2	-	0/9/10/12	0/2/2/2
2	A1MCG	D	6	2	-	2/9/10/12	0/2/2/2
2	A1MCG	B	8	2	-	2/9/10/12	0/2/2/2
2	A1MCG	B	6	2	-	2/9/10/12	0/2/2/2
2	A1MCG	D	9	2	-	2/9/10/12	0/2/2/2

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	9	A1MCG	C14-N13	5.06	1.46	1.36
2	D	9	A1MCG	C14-N13	4.84	1.46	1.36
2	D	6	A1MCG	C14-N13	4.63	1.46	1.36
2	B	8	A1MCG	C14-N13	4.16	1.45	1.36
2	D	8	A1MCG	C14-N13	4.01	1.44	1.36
2	B	6	A1MCG	C14-N13	3.97	1.44	1.36
2	D	6	A1MCG	CZ-N13	3.22	1.47	1.40
2	D	9	A1MCG	CZ-N13	2.95	1.47	1.40
2	D	8	A1MCG	C14-N19	2.58	1.38	1.34
2	B	6	A1MCG	C14-N19	2.51	1.38	1.34
2	B	9	A1MCG	CZ-N13	2.49	1.46	1.40
2	D	9	A1MCG	C14-N19	2.43	1.38	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	6	A1MCG	CZ-N13	2.37	1.46	1.40
2	D	6	A1MCG	C14-N19	2.25	1.37	1.34
2	B	9	A1MCG	O-C	2.06	1.28	1.19

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	9	A1MCG	C16-N15-C14	8.56	123.04	115.45
2	D	6	A1MCG	C16-N15-C14	8.43	122.92	115.45
2	D	9	A1MCG	C16-N15-C14	7.41	122.02	115.45
2	D	6	A1MCG	N19-C14-N15	-7.38	118.66	126.43
2	D	9	A1MCG	N19-C14-N15	-6.95	119.11	126.43
2	B	9	A1MCG	N19-C14-N15	-6.74	119.33	126.43
2	B	6	A1MCG	C16-N15-C14	5.92	120.70	115.45
2	B	8	A1MCG	N19-C14-N15	-4.67	121.51	126.43
2	D	8	A1MCG	C16-N15-C14	4.63	119.56	115.45
2	B	6	A1MCG	N19-C14-N15	-4.17	122.03	126.43
2	D	6	A1MCG	CZ-N13-C14	-3.84	118.09	129.23
2	B	8	A1MCG	C16-N15-C14	3.77	118.80	115.45
2	D	8	A1MCG	N19-C14-N15	-3.71	122.52	126.43
2	B	8	A1MCG	CD2-CE2-CZ	-3.64	116.09	120.30
2	B	8	A1MCG	C18-N19-C14	3.56	118.61	115.45
2	B	6	A1MCG	C18-C17-C16	3.55	121.75	116.67
2	D	8	A1MCG	C18-C17-C16	3.46	121.62	116.67
2	D	8	A1MCG	CG-CB-CA	3.20	120.58	114.10
2	D	9	A1MCG	C18-N19-C14	3.13	118.23	115.45
2	B	9	A1MCG	C17-C16-N15	-2.94	118.62	123.43
2	D	9	A1MCG	CB-CA-C	-2.90	106.04	111.47
2	B	6	A1MCG	C17-C16-N15	-2.89	118.70	123.43
2	B	8	A1MCG	C18-C17-C16	2.79	120.67	116.67
2	D	6	A1MCG	CB-CA-C	-2.75	106.31	111.47
2	B	6	A1MCG	CZ-N13-C14	-2.72	121.34	129.23
2	D	6	A1MCG	C17-C16-N15	-2.60	119.19	123.43
2	B	8	A1MCG	CE2-CZ-CE1	2.58	122.56	119.03
2	B	8	A1MCG	CB-CA-C	-2.54	106.70	111.47
2	B	8	A1MCG	CB-CG-CD1	-2.54	115.86	120.91
2	B	8	A1MCG	C17-C18-N19	-2.50	119.35	123.43
2	B	9	A1MCG	C18-C17-C16	2.43	120.15	116.67
2	D	9	A1MCG	C18-C17-C16	2.39	120.10	116.67
2	D	6	A1MCG	C18-N19-C14	2.36	117.55	115.45
2	D	8	A1MCG	C17-C16-N15	-2.31	119.65	123.43
2	D	9	A1MCG	C17-C16-N15	-2.25	119.75	123.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	9	A1MCG	CZ-N13-C14	-2.22	122.78	129.23
2	D	8	A1MCG	CB-CA-C	-2.15	107.44	111.47
2	D	8	A1MCG	C17-C18-N19	-2.05	120.07	123.43
2	B	8	A1MCG	CG-CB-CA	2.02	118.19	114.10

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	9	A1MCG	CA-CB-CG-CD1
2	B	9	A1MCG	CA-CB-CG-CD2
2	D	9	A1MCG	CA-CB-CG-CD1
2	D	9	A1MCG	CA-CB-CG-CD2
2	B	6	A1MCG	CE1-CZ-N13-C14
2	D	6	A1MCG	CE1-CZ-N13-C14
2	D	6	A1MCG	CE2-CZ-N13-C14
2	B	6	A1MCG	CE2-CZ-N13-C14
2	B	8	A1MCG	CA-CB-CG-CD1
2	B	8	A1MCG	CA-CB-CG-CD2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	9	A1MCG	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PRO	A	502	-	8,8,8	4.15	3 (37%)	10,10,10	3.04	5 (50%)
5	PEG	A	503	-	6,6,6	0.41	0	5,5,5	0.45	0
4	PRO	C	502	-	8,8,8	2.36	2 (25%)	10,10,10	2.23	3 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PRO	A	502	-	-	2/4/11/11	0/1/1/1
5	PEG	A	503	-	-	4/4/4/4	-
4	PRO	C	502	-	-	3/4/11/11	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	502	PRO	CG-CB	10.59	1.95	1.51
4	C	502	PRO	CG-CB	5.46	1.74	1.51
4	A	502	PRO	O-C	3.40	1.32	1.22
4	A	502	PRO	CA-C	3.09	1.60	1.52
4	C	502	PRO	OXT-C	-2.35	1.22	1.30

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	502	PRO	O-C-CA	6.39	143.22	122.26
4	C	502	PRO	OXT-C-O	-4.50	113.88	124.09
4	A	502	PRO	OXT-C-O	-4.31	114.30	124.09
4	C	502	PRO	OXT-C-CA	3.70	125.70	113.40
4	C	502	PRO	C-CA-N	3.35	119.93	106.73
4	A	502	PRO	OXT-C-CA	-3.29	102.47	113.40
4	A	502	PRO	C-CA-N	3.24	119.51	106.73
4	A	502	PRO	CG-CD-N	-2.12	99.47	105.69

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	502	PRO	O-C-CA-N

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
4	A	502	PRO	OXT-C-CA-N
4	C	502	PRO	O-C-CA-N
4	C	502	PRO	OXT-C-CA-N
5	A	503	PEG	O2-C3-C4-O4
5	A	503	PEG	C1-C2-O2-C3
5	A	503	PEG	O1-C1-C2-O2
5	A	503	PEG	C4-C3-O2-C2
4	C	502	PRO	O-C-CA-CB

There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	502	PRO	5	0
4	C	502	PRO	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	349/359 (97%)	0.20	22 (6%)	27 29	12, 34, 62, 80	4 (1%)
1	C	344/359 (95%)	0.24	14 (4%)	42 45	11, 41, 68, 95	7 (2%)
2	B	7/17 (41%)	1.42	3 (42%)	1 0	40, 57, 68, 77	0
2	D	7/17 (41%)	2.21	3 (42%)	1 0	54, 68, 76, 85	0
All	All	707/752 (94%)	0.25	42 (5%)	29 31	11, 38, 68, 95	11 (1%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	373	LEU	5.8
2	D	10	TYR	5.2
1	A	372	PRO	5.1
1	A	374	ALA	4.8
1	A	375	THR	4.4
1	A	376	ILE	3.6
1	A	124	ASP	3.4
1	A	47	GLY	3.3
1	C	297	LYS	3.2
2	D	14	SER	3.2
1	C	298	ALA	3.1
2	B	10	TYR	3.0
1	A	93	PHE	3.0
1	C	71	TYR	3.0
1	A	48	PRO	2.9
1	A	371	PRO	2.9
1	A	35	SER	2.8
1	C	296	ILE	2.8
1	A	67	PHE	2.7
1	A	64	ASN	2.6
1	C	300	PRO	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	124	ASP	2.5
1	A	69	VAL	2.5
1	A	383	ARG	2.4
1	C	36	LYS	2.4
1	A	121	GLU	2.4
1	A	50	ARG	2.3
1	C	299	HIS	2.2
1	C	358	ALA	2.2
2	B	5	ASP	2.2
1	A	65	GLY	2.2
1	C	302	THR	2.2
2	B	14	SER	2.1
1	A	123	LYS	2.1
1	C	290	GLU	2.1
1	C	119	SER	2.1
1	A	353	GLY	2.1
1	A	46	GLN	2.1
1	A	77	ASP	2.1
1	C	91	LYS	2.0
1	C	77	ASP	2.0
2	D	5	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	A1MCG	D	9	18/19	0.88	0.13	63,69,81,82	0
2	A1MCG	D	6	18/19	0.90	0.13	56,62,69,70	0
2	A1MCG	B	9	18/19	0.91	0.11	47,61,76,76	0
2	A1MCG	D	8	18/19	0.93	0.12	55,56,61,61	0
2	A1MCG	B	6	18/19	0.94	0.09	43,48,54,57	0
2	A1MCG	B	8	18/19	0.95	0.09	40,42,48,49	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PRO	A	502	8/8	0.91	0.15	27,30,48,64	0
4	PRO	C	502	8/8	0.91	0.14	27,31,46,58	0
5	PEG	A	503	7/7	0.92	0.12	54,56,64,65	0
3	CL	C	501	1/1	0.98	0.12	39,39,39,39	0
3	CL	A	501	1/1	0.99	0.08	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.