



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 3, 2026 – 01:12 AM JST

PDB ID : 9X5I / pdb_00009x5i
Title : Crystal structure of Medicago truncatula DELLA1-GRAS domain
Authors : Wan, L.H.
Deposited on : 2025-10-13
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	 3% 84% 7% 9%
1	B	387	 7% 81% 9% 10%
1	C	387	 3% 81% 9% 9%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8805 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DELLA protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	352	2743	1751	481	498	13	0	0	0
1	B	347	2711	1735	474	489	13	0	1	0
1	C	351	2742	1751	480	498	13	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	386	SER	CYS	engineered mutation	UNP A0A396IUP1
B	386	SER	CYS	engineered mutation	UNP A0A396IUP1
C	386	SER	CYS	engineered mutation	UNP A0A396IUP1

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	201	Total 201	O 201	0	0
2	B	199	Total 199	O 199	0	0
2	C	209	Total 209	O 209	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	91.44Å 91.44Å 229.08Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.18 – 2.00 38.18 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.7 (38.18-2.00) 97.8 (38.18-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.76 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.181 , 0.209 0.181 , 0.209	Depositor DCC
R_{free} test set	3553 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å ²)	23.5	Xtrriage
Anisotropy	0.068	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8805	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/2802	0.46	0/3793
1	B	0.29	0/2773	0.46	0/3754
1	C	0.35	0/2801	0.53	0/3792
All	All	0.31	0/8376	0.49	0/11339

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2743	0	2733	25	0
1	B	2711	0	2712	32	0
1	C	2742	0	2731	29	0
2	A	201	0	0	6	0
2	B	199	0	0	8	0
2	C	209	0	0	3	0
All	All	8805	0	8176	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:269:GLY:O	1:C:270:ASN:ND2	2.09	0.84
1:B:420:PRO:HA	2:B:658:HOH:O	1.81	0.79
1:A:547:GLN:HE21	1:B:220:MET:HG3	1.51	0.75
1:A:554:LEU:HB3	1:B:252:MET:HG3	1.68	0.74
1:B:538:VAL:HG22	1:B:586:LYS:HE3	1.72	0.70
1:C:444:GLU:HG2	1:C:571:MET:HE1	1.72	0.70
1:B:240:HIS:HA	1:B:243:LEU:HD12	1.75	0.69
1:B:211:THR:N	2:B:605:HOH:O	2.28	0.67
1:B:389:ILE:HD12	2:B:701:HOH:O	1.94	0.67
1:B:554:LEU:HD22	1:C:252:MET:HA	1.78	0.66
1:C:346:ARG:HH22	1:C:398:GLU:HG3	1.62	0.65
1:B:265:ARG:NH2	1:B:506:CYS:SG	2.70	0.64
1:C:346:ARG:NH2	1:C:398:GLU:HG3	2.13	0.64
1:B:286:HIS:HB2	1:B:496:MET:HE3	1.82	0.62
1:C:356:ASP:N	1:C:356:ASP:OD1	2.31	0.61
1:A:547:GLN:NE2	1:B:220:MET:HG3	2.15	0.60
1:B:554:LEU:HB3	1:C:252:MET:HB2	1.84	0.60
1:C:354:GLN:HB3	1:C:356:ASP:OD1	2.01	0.59
1:A:286:HIS:ND1	1:A:493:ASP:OD1	2.34	0.59
1:A:211:THR:N	2:A:605:HOH:O	2.36	0.59
1:C:444:GLU:CG	1:C:571:MET:HE1	2.32	0.59
1:C:571:MET:HE3	1:C:581:ALA:HB2	1.83	0.58
1:B:326:MET:HE3	2:B:647:HOH:O	2.04	0.56
1:C:527:ARG:NH2	1:C:567:ASN:HB2	2.20	0.56
1:A:224:GLU:O	1:A:228:GLN:HG2	2.06	0.55
1:B:373:GLN:CD	2:B:603:HOH:O	2.51	0.53
1:A:536:GLU:OE2	1:A:586:LYS:HE2	2.10	0.52
1:A:436:LYS:HE3	2:A:703:HOH:O	2.09	0.52
1:C:214:ARG:HD2	1:C:244:LEU:HD11	1.92	0.52
1:C:536:GLU:OE2	1:C:588:PRO:HG3	2.10	0.52
1:B:546:LYS:HE3	1:C:458:GLU:OE1	2.09	0.52
1:A:538:VAL:HG22	1:A:586:LYS:HG2	1.93	0.51
1:C:588:PRO:HD2	2:C:687:HOH:O	2.10	0.50
1:B:243:LEU:O	1:B:246:SER:OG	2.26	0.50
1:B:244:LEU:HA	1:B:247:LEU:HD12	1.94	0.50
1:B:263:LEU:O	1:B:267:ILE:HG12	2.12	0.50
1:C:285:MET:HE1	1:C:326:MET:HE2	1.93	0.49
1:B:234:ALA:O	1:B:238:VAL:HG23	2.13	0.49
1:B:215:LEU:HD13	1:B:252:MET:SD	2.53	0.49
1:B:238:VAL:HG21	1:B:267:ILE:HG13	1.95	0.49
1:B:455:ARG:NH2	2:B:604:HOH:O	2.44	0.48
1:B:330:ALA:HB3	2:B:776:HOH:O	2.12	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:376:GLY:HA2	2:A:748:HOH:O	2.13	0.47
1:C:333:GLN:O	1:C:337:LEU:HG	2.15	0.47
1:A:425:LYS:NZ	2:A:611:HOH:O	2.47	0.47
1:A:366:TRP:O	1:A:370:GLN:HG3	2.15	0.46
1:B:211:THR:CA	2:B:605:HOH:O	2.64	0.46
1:A:554:LEU:HD13	1:B:252:MET:HG3	1.98	0.46
1:B:437:ILE:HD11	1:B:584:ALA:HB1	1.97	0.46
1:A:338:ARG:HD2	2:A:666:HOH:O	2.15	0.46
1:A:404:ALA:HB1	1:A:436:LYS:HG3	1.98	0.45
1:C:317:VAL:HG11	1:C:328:TRP:CH2	2.51	0.45
1:C:282:ILE:HG23	1:C:286:HIS:CE1	2.51	0.45
1:C:350:ILE:HD13	1:C:392:LEU:HD11	1.98	0.45
1:B:286:HIS:ND1	1:B:496:MET:HB2	2.32	0.45
1:C:282:ILE:HD13	1:C:282:ILE:HA	1.92	0.44
1:C:283:LEU:HD23	1:C:496:MET:HE3	1.99	0.43
1:A:257:SER:O	1:A:261:GLN:HG3	2.17	0.43
1:A:545:PHE:CD1	1:A:565:GLU:HG3	2.53	0.43
1:A:571:MET:HE3	1:A:571:MET:HB3	1.91	0.43
1:A:588:PRO:O	1:A:589:GLN:HG3	2.19	0.43
1:C:279:PHE:HE1	1:C:496:MET:HE1	1.84	0.43
1:A:337:LEU:HD13	1:B:237:LEU:HD21	2.01	0.43
1:B:248:GLN:HG2	1:B:252:MET:SD	2.59	0.42
1:A:261:GLN:O	1:A:265:ARG:HG3	2.18	0.42
1:A:540:LEU:HD23	1:A:540:LEU:HA	1.81	0.42
1:A:536:GLU:OE2	1:A:588:PRO:HA	2.19	0.42
1:A:228:GLN:HG3	1:C:339:PRO:HG2	2.02	0.41
1:C:424:GLU:HG3	2:C:807:HOH:O	2.19	0.41
1:C:466:LEU:HD23	1:C:466:LEU:HA	1.90	0.41
1:C:539:HIS:HB3	2:C:673:HOH:O	2.20	0.41
1:B:237:LEU:O	1:B:241:ILE:HG13	2.20	0.41
1:C:527:ARG:NH2	1:C:566:ASN:O	2.54	0.41
1:A:403:GLU:OE1	2:A:601:HOH:O	2.22	0.41
1:C:436:LYS:HB2	1:C:436:LYS:HE2	1.90	0.41
1:B:404:ALA:HB1	1:B:436:LYS:HG3	2.02	0.40
1:B:571:MET:HE3	1:B:571:MET:HB3	2.00	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	344/387 (89%)	338 (98%)	6 (2%)	0	100	100
1	B	340/387 (88%)	335 (98%)	5 (2%)	0	100	100
1	C	343/387 (89%)	338 (98%)	5 (2%)	0	100	100
All	All	1027/1161 (88%)	1011 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/319 (91%)	289 (100%)	0	100	100
1	B	286/319 (90%)	286 (100%)	0	100	100
1	C	289/319 (91%)	289 (100%)	0	100	100
All	All	864/957 (90%)	864 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	261	GLN
1	A	434	ASN
1	B	333	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	378	GLN
1	B	387	ASN
1	B	539	HIS
1	C	327	GLN
1	C	370	GLN
1	C	408	ASN
1	C	434	ASN
1	C	589	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	352/387 (90%)	-0.12	10 (2%) 55 54	15, 28, 51, 84	0
1	B	347/387 (89%)	0.10	27 (7%) 19 18	14, 30, 70, 84	1 (0%)
1	C	351/387 (90%)	-0.04	10 (2%) 55 54	14, 29, 60, 100	0
All	All	1050/1161 (90%)	-0.02	47 (4%) 38 37	14, 29, 62, 100	1 (0%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	555	PHE	4.4
1	C	553	ALA	4.1
1	B	243	LEU	4.1
1	B	555	PHE	3.9
1	B	252	MET	3.8
1	A	554	LEU	3.8
1	A	553	ALA	3.7
1	A	493	ASP	3.6
1	C	555	PHE	3.6
1	B	241	ILE	3.5
1	B	267	ILE	3.4
1	B	249	THR	3.3
1	A	494	LEU	3.2
1	A	232	LYS	3.1
1	B	494	LEU	3.0
1	C	269	GLY	2.9
1	B	269	GLY	2.9
1	B	244	LEU	2.8
1	B	225	ALA	2.7
1	B	553	ALA	2.7
1	B	233	LEU	2.6
1	B	276	ASP	2.6
1	C	358	THR	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	252	MET	2.6
1	B	554	LEU	2.5
1	B	366	TRP	2.5
1	B	215	LEU	2.4
1	B	247	LEU	2.4
1	C	366	TRP	2.3
1	B	250	GLY	2.3
1	B	226	ILE	2.3
1	B	238	VAL	2.3
1	B	236	ALA	2.3
1	B	237	LEU	2.3
1	B	375	ILE	2.3
1	C	554	LEU	2.2
1	A	366	TRP	2.2
1	C	270	ASN	2.2
1	C	356	ASP	2.2
1	B	231	LEU	2.2
1	B	246	SER	2.2
1	A	270	ASN	2.2
1	A	589	GLN	2.1
1	B	420	PRO	2.1
1	A	491	SER	2.1
1	C	240	HIS	2.0
1	B	211	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.