Full wwPDB X-ray Structure Validation Report ⓘ

Nov 16, 2026 – 09:04 PM JST

PDB ID	:	9XQ2 / pdb_00009xq2
Title	:	Crystal structure of SARS-Cov-2 main protease S144A mutant in complex with CDD1845
Authors	:	Li, W.W.; Zeng, P.; Guo, L.; Zhou, X.L.; Li, J.
Deposited on	:	2025-11-17
Resolution	:	1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	6% 88% 8% .
1	B	306	15% 86% 10% . .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4809 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3C-like proteinase nsp5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	295	Total	C	N	O	S	0	0	0
			2247	1425	378	423	21			
1	B	296	Total	C	N	O	S	0	0	0
			2251	1431	377	422	21			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	144	ALA	SER	engineered mutation	UNP P0DTD1
B	144	ALA	SER	engineered mutation	UNP P0DTD1

- Molecule 2 is (2P)-2-(isoquinolin-4-yl)-1-[4-(methylamino)-4-oxobutyl]-N-[(1S)-1-(naphthalen-2-yl)ethyl]-1H-benzimidazole-7-carboxamide (CCD ID: O5F) (formula: $C_{34}H_{31}N_5O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			41	34	5	2		
2	B	1	Total	C	N	O	0	0
			41	34	5	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	142	Total	O	0	0
			142	142		
3	B	87	Total	O	0	0
			87	87		


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 3C-like proteinase nsp5



• Molecule 1: 3C-like proteinase nsp5



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	68.00Å 101.59Å 105.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.09 – 1.95 57.09 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.8 (57.09-1.95) 99.8 (57.09-1.95)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.70 (at 1.95Å)	Xtriage
Refinement program	PHENIX 1.12_2829, PHENIX 1.12_2829	Depositor
R, R_{free}	0.204 , 0.244 0.206 , 0.247	Depositor DCC
R_{free} test set	2720 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å^2)	25.7	Xtriage
Anisotropy	0.596	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.38 , 37.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.014 for -h,l,k	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	4809	wwPDB-VP
Average B, all atoms (Å^2)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: O5F

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.32	0/2297	0.56	0/3129
1	B	0.32	0/2302	0.54	0/3136
All	All	0.32	0/4599	0.55	0/6265

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2247	0	2168	19	0
1	B	2251	0	2167	26	0
2	A	41	0	0	0	0
2	B	41	0	0	2	0
3	A	142	0	0	2	0
3	B	87	0	0	3	0
All	All	4809	0	4335	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:154:TYR:O	1:B:155:ASP:OD2	1.80	0.99
1:B:276:MET:HE3	1:B:281:ILE:HG12	1.65	0.76
1:A:102:LYS:NZ	1:A:156:CYS:SG	2.65	0.69
1:B:142:ASN:HA	2:B:401:O5F:C27	2.27	0.65
1:B:3:PHE:N	3:B:503:HOH:O	2.33	0.61
1:A:5:LYS:HG3	1:A:291:PHE:CZ	2.36	0.61
1:A:109:GLY:HA2	1:A:200:ILE:HD13	1.83	0.61
1:B:221:ASN:HD21	1:B:223:PHE:HB2	1.67	0.58
1:A:167:LEU:HD21	1:A:173:ALA:HB2	1.87	0.56
1:B:188:ARG:NE	3:B:506:HOH:O	2.39	0.55
1:A:128:CYS:SG	1:A:136:ILE:HG23	2.49	0.53
1:A:231:ASN:HB3	1:A:235:MET:HE2	1.90	0.53
1:A:126:TYR:CD1	1:B:4:ARG:HD2	2.45	0.52
1:B:3:PHE:N	3:B:508:HOH:O	2.44	0.51
1:B:276:MET:O	1:B:279:ARG:HG3	2.11	0.51
1:B:233:VAL:HG21	1:B:269:LYS:HD2	1.94	0.50
1:B:143:GLY:HA2	2:B:401:O5F:C16	2.44	0.46
1:A:108:PRO:HB3	1:A:132:PRO:HA	1.96	0.46
1:B:113:SER:O	1:B:149:GLY:HA2	2.16	0.46
1:A:233:VAL:HG21	1:A:269:LYS:HE3	1.98	0.45
1:B:45:THR:OG1	1:B:47:GLU:HG3	2.15	0.45
1:A:126:TYR:HD2	1:B:6:MET:HE3	1.81	0.45
1:B:219:PHE:CD2	1:B:281:ILE:HD12	2.52	0.45
1:A:207:TRP:CH2	1:A:281:ILE:HB	2.52	0.45
1:A:5:LYS:HD3	3:A:515:HOH:O	2.17	0.45
1:A:288:GLU:HG2	1:A:291:PHE:CE2	2.52	0.44
1:A:288:GLU:HG2	1:A:291:PHE:CD2	2.53	0.44
1:A:124:GLY:HA3	1:B:6:MET:HE2	1.99	0.44
1:B:102:LYS:HB2	1:B:102:LYS:HE2	1.78	0.43
1:B:216:ASP:OD2	1:B:281:ILE:O	2.37	0.43
1:B:221:ASN:ND2	1:B:223:PHE:H	2.16	0.42
1:B:86:VAL:HG13	1:B:179:GLY:HA2	2.01	0.42
1:B:15:GLY:O	1:B:97:LYS:HE2	2.20	0.42
1:A:207:TRP:CE2	1:A:288:GLU:HB3	2.55	0.42
1:A:56:ASP:OD1	3:A:501:HOH:O	2.22	0.42
1:A:86:VAL:HG13	1:A:179:GLY:HA2	2.03	0.41
1:A:188:ARG:HG2	1:A:190:THR:HG23	2.03	0.41
1:B:207:TRP:CH2	1:B:281:ILE:HB	2.56	0.40
1:B:108:PRO:HG3	1:B:134:PHE:CE1	2.56	0.40
1:B:154:TYR:O	1:B:155:ASP:CG	2.59	0.40
1:B:220:LEU:HD23	1:B:220:LEU:HA	1.91	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:246:HIS:HA	1:B:249:ILE:HG22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	293/306 (96%)	287 (98%)	6 (2%)	0	100 100
1	B	294/306 (96%)	280 (95%)	10 (3%)	4 (1%)	9 3
All	All	587/612 (96%)	567 (97%)	16 (3%)	4 (1%)	18 10

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	142	ASN
1	B	46	SER
1	B	154	TYR
1	B	155	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	244/262 (93%)	244 (100%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	243/262 (93%)	243 (100%)	0	100 100
All	All	487/524 (93%)	487 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
1	A	107	GLN
1	A	277	ASN
1	B	69	GLN
1	B	221	ASN
1	B	274	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
2	O5F	B	401	-	46,46,46	5.43	32 (69%)	61,65,65	1.68	11 (18%)
2	O5F	A	401	-	46,46,46	5.28	32 (69%)	61,65,65	1.83	12 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	O5F	B	401	-	-	0/24/24/24	0/6/6/6
2	O5F	A	401	-	-	0/24/24/24	0/6/6/6

All (64) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	O5F	C13-C18	12.93	1.58	1.40
2	A	401	O5F	C13-C18	11.85	1.56	1.40
2	B	401	O5F	C16-C17	11.32	1.58	1.39
2	A	401	O5F	C16-C17	10.48	1.57	1.39
2	A	401	O5F	C32-N3	9.25	1.48	1.32
2	B	401	O5F	C32-N3	9.18	1.48	1.32
2	B	401	O5F	C3-C2	8.73	1.53	1.39
2	A	401	O5F	C3-C2	8.59	1.52	1.39
2	B	401	O5F	C4-C3	7.74	1.52	1.36
2	A	401	O5F	C4-C3	7.65	1.52	1.36
2	B	401	O5F	C7-C6	7.40	1.53	1.36
2	B	401	O5F	C11-C2	7.40	1.52	1.37
2	A	401	O5F	C33-C25	7.36	1.52	1.39
2	B	401	O5F	C8-C9	7.32	1.53	1.36
2	A	401	O5F	C7-C6	7.28	1.53	1.36
2	A	401	O5F	C8-C9	7.23	1.53	1.36
2	B	401	O5F	C28-C27	7.15	1.53	1.36
2	A	401	O5F	C11-C2	7.15	1.51	1.37
2	A	401	O5F	C29-C30	7.00	1.52	1.36
2	B	401	O5F	C29-C30	7.00	1.52	1.36
2	B	401	O5F	C33-C25	6.98	1.52	1.39
2	A	401	O5F	C28-C27	6.98	1.52	1.36
2	B	401	O5F	C33-N3	6.76	1.48	1.34
2	B	401	O5F	C27-C26	6.46	1.55	1.42
2	B	401	O5F	C14-C13	6.22	1.49	1.39
2	B	401	O5F	C32-C31	6.21	1.54	1.41
2	A	401	O5F	C33-N3	6.15	1.47	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	O5F	C27-C26	6.15	1.54	1.42
2	A	401	O5F	C8-C7	6.13	1.54	1.38
2	B	401	O5F	C8-C7	6.06	1.54	1.38
2	A	401	O5F	C32-C31	5.99	1.54	1.41
2	A	401	O5F	C14-C13	5.91	1.49	1.39
2	B	401	O5F	C29-C28	5.81	1.53	1.38
2	A	401	O5F	C29-C28	5.78	1.53	1.38
2	B	401	O5F	C15-C16	5.56	1.50	1.38
2	B	401	O5F	C11-C10	5.55	1.54	1.42
2	B	401	O5F	C4-C5	5.49	1.55	1.41
2	B	401	O5F	C30-C31	5.49	1.55	1.41
2	A	401	O5F	C4-C5	5.44	1.55	1.41
2	A	401	O5F	C30-C31	5.38	1.55	1.41
2	A	401	O5F	C6-C5	5.24	1.54	1.41
2	A	401	O5F	C11-C10	5.16	1.53	1.42
2	B	401	O5F	C6-C5	5.16	1.54	1.41
2	B	401	O5F	C9-C10	4.98	1.54	1.41
2	A	401	O5F	C9-C10	4.94	1.53	1.41
2	A	401	O5F	C15-C16	4.82	1.49	1.38
2	B	401	O5F	C12-N	4.71	1.44	1.34
2	A	401	O5F	C12-N	4.61	1.44	1.34
2	B	401	O5F	C25-C26	4.51	1.51	1.43
2	A	401	O5F	C25-C26	4.50	1.51	1.43
2	A	401	O5F	C17-N4	-3.96	1.31	1.39
2	B	401	O5F	C15-C14	3.85	1.47	1.38
2	A	401	O5F	C15-C14	3.78	1.46	1.38
2	A	401	O5F	C10-C5	3.60	1.50	1.42
2	A	401	O5F	C18-N1	-3.53	1.33	1.40
2	B	401	O5F	C10-C5	3.48	1.50	1.42
2	B	401	O5F	C18-N1	-3.28	1.33	1.40
2	B	401	O5F	C17-N4	-2.72	1.34	1.39
2	B	401	O5F	C22-N2	2.54	1.46	1.34
2	A	401	O5F	C22-N2	2.33	1.45	1.34
2	A	401	O5F	C26-C31	2.31	1.47	1.43
2	B	401	O5F	C26-C31	2.30	1.47	1.43
2	A	401	O5F	C24-N1	-2.23	1.35	1.38
2	B	401	O5F	C24-N4	2.21	1.36	1.32

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	401	O5F	C17-C18-N1	6.61	108.42	104.71

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	401	O5F	C32-N3-C33	5.70	121.70	116.87
2	B	401	O5F	C17-C18-N1	5.37	107.72	104.71
2	B	401	O5F	C32-N3-C33	5.27	121.34	116.87
2	A	401	O5F	C25-C33-N3	-4.57	117.41	124.17
2	B	401	O5F	C25-C33-N3	-3.90	118.40	124.17
2	B	401	O5F	C25-C24-N1	2.84	128.30	123.64
2	A	401	O5F	C20-C19-N1	-2.84	106.40	112.44
2	A	401	O5F	C3-C2-C11	2.58	121.93	118.60
2	B	401	O5F	C20-C21-C22	-2.49	106.27	113.26
2	B	401	O5F	C32-C31-C26	2.47	120.15	117.78
2	B	401	O5F	C30-C31-C32	-2.36	118.40	122.63
2	A	401	O5F	C21-C22-N2	2.34	119.54	116.29
2	B	401	O5F	C19-N1-C24	-2.31	124.38	128.91
2	B	401	O5F	C20-C19-N1	-2.29	107.56	112.44
2	A	401	O5F	C13-C18-C17	-2.28	118.53	121.25
2	A	401	O5F	C9-C10-C11	-2.14	118.24	122.02
2	B	401	O5F	C27-C26-C31	2.13	120.64	117.89
2	A	401	O5F	C30-C31-C32	-2.11	118.86	122.63
2	A	401	O5F	C20-C21-C22	-2.05	107.50	113.26
2	A	401	O5F	C27-C26-C31	2.03	120.52	117.89
2	B	401	O5F	C27-C26-C25	-2.01	120.22	123.31
2	A	401	O5F	C19-N1-C24	-2.00	124.98	128.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.


1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	O5F	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.


Ligand O5F B 401




Bond lengths



Bond angles




Torsions




Rings


Ligand O5F A 401




Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	295/306 (96%)	0.34	19 (6%) 25 29	16, 26, 40, 46	0
1	B	296/306 (96%)	0.89	47 (15%) 5 5	17, 32, 51, 61	0
All	All	591/612 (96%)	0.62	66 (11%) 10 12	16, 28, 48, 61	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	142	ASN	6.8
1	B	281	ILE	5.3
1	B	215	GLY	4.9
1	B	223	PHE	4.7
1	B	224	THR	4.4
1	A	291	PHE	4.2
1	B	24	THR	4.1
1	B	255	ALA	3.7
1	A	154	TYR	3.7
1	B	154	TYR	3.6
1	A	72	ASN	3.5
1	A	282	LEU	3.6
1	B	294	PHE	3.4
1	B	221	ASN	3.4
1	B	92	ASP	3.4
1	B	153	ASP	3.4
1	B	59	ILE	3.3
1	B	298	ARG	3.3
1	B	72	ASN	3.3
1	B	282	LEU	3.3
1	B	3	PHE	3.2
1	B	47	GLU	3.2
1	B	297	VAL	3.2
1	B	46	SER	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	155	ASP	3.0
1	B	249	ILE	3.0
1	A	216	ASP	3.0
1	B	278	GLY	2.9
1	B	222	ARG	2.9
1	B	245	ASP	2.8
1	A	153	ASP	2.8
1	B	64	HIS	2.7
1	A	294	PHE	2.7
1	B	218	TRP	2.6
1	B	232	LEU	2.6
1	A	298	ARG	2.6
1	B	214	ASN	2.6
1	A	191	ALA	2.6
1	A	283	GLY	2.5
1	B	23	GLY	2.5
1	A	92	ASP	2.4
1	B	276	MET	2.4
1	B	229	ASP	2.4
1	B	77	VAL	2.4
1	A	215	GLY	2.4
1	A	136	ILE	2.3
1	A	128	CYS	2.3
1	B	216	ASP	2.3
1	B	270	GLU	2.3
1	B	233	VAL	2.3
1	A	214	ASN	2.2
1	B	67	LEU	2.2
1	A	77	VAL	2.2
1	B	273	GLN	2.2
1	A	155	ASP	2.2
1	B	213	ILE	2.1
1	B	73	VAL	2.1
1	B	196	THR	2.1
1	A	4	ARG	2.1
1	A	67	LEU	2.1
1	B	60	ARG	2.1
1	B	81	SER	2.1
1	B	263	ASP	2.1
1	B	217	ARG	2.1
1	B	220	LEU	2.0
1	B	230	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	O5F	B	401	41/41	0.92	0.11	25,30,34,39	0
2	O5F	A	401	41/41	0.95	0.08	16,21,26,29	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around O5F B 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)








Electron density around O5F A 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.