Full wwPDB X-ray Structure Validation Report (i)

Nov 16, 2026 – 08:56 PM JST

PDB ID	:	9XQP / pdb_00009xqp
Title	:	Crystal structure of FGFR4 kinase domain with inhibitor F-30
Authors	:	Zhang, L.; Chen, X.
Deposited on	:	2025-11-18
Resolution	:	2.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6062 (2.44-2.40)
Clashscore	190562	6562 (2.44-2.40)
Ramachandran outliers	187476	6481 (2.44-2.40)
Sidechain outliers	187428	6482 (2.44-2.40)
RSRZ outliers	180081	6066 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	5%, 79%, 14%, • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	802	-	-	X	-
3	SO4	A	803	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2425 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Fibroblast growth factor receptor 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	294	Total	C	N	O	S	0	0	0
			2313	1478	403	416	16			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	443	GLY	-	expression tag	UNP P22455
A	444	PRO	-	expression tag	UNP P22455
A	664	GLU	ARG	conflict	UNP P22455

- Molecule 2 is 3-(2-chloranylethanoylamino)- {N}-[6-(3,5-dimethoxyphenyl)-1 {H}-indazol-3 -yl]benzamide (CCD ID: A1E09) (formula: $C_{24}H_{21}ClN_4O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			32	24	4	4		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	60	Total O 60 60	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Fibroblast growth factor receptor 4



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	42.36Å 57.56Å 59.76Å 90.00° 91.60° 90.00°	Depositor
Resolution (Å)	42.35 – 2.43 42.35 – 2.43	Depositor EDS
% Data completeness (in resolution range)	95.8 (42.35-2.43) 95.7 (42.35-2.43)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.99 (at 2.42Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.216 , 0.278 0.218 , 0.278	Depositor DCC
R_{free} test set	475 reflections (4.31%)	wwPDB-VP
Wilson B-factor (Å^2)	30.8	Xtriage
Anisotropy	0.448	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 35.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k 0.019 for -h,-l,-k 0.046 for h,-k,-l	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	2425	wwPDB-VP
Average B, all atoms (Å^2)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.91% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1E09, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.42	0/2371	0.86	2/3217 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	507	ASP	CB-CA-C	6.09	119.83	109.53
1	A	560	GLU	CB-CG-CD	5.02	121.13	112.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	611	ARG	Sidechain
1	A	616	ARG	Sidechain
1	A	635	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2313	0	2306	33	0
2	A	32	0	0	0	0
3	A	20	0	0	6	0
4	A	60	0	0	5	0
All	All	2425	0	2306	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:526:LEU:O	1:A:606:ARG:NH1	1.98	0.96
1:A:464:ARG:NH1	1:A:545:PRO:O	2.10	0.84
1:A:456:LEU:HG	1:A:461:GLU:HG3	1.72	0.72
1:A:506:LYS:HZ2	1:A:506:LYS:HB3	1.58	0.68
1:A:663:ASP:OD2	4:A:902:HOH:O	2.10	0.67
1:A:540:CYS:HB2	1:A:547:TYR:HB2	1.77	0.66
1:A:464:ARG:NH2	1:A:542:GLN:O	2.29	0.66
1:A:456:LEU:HD12	1:A:542:GLN:HG3	1.81	0.63
1:A:503:LYS:NZ	4:A:901:HOH:O	2.09	0.62
1:A:646:THR:HG23	1:A:648:ASN:H	1.66	0.59
1:A:646:THR:HG21	3:A:802:SO4:S	2.43	0.59
1:A:646:THR:HG22	1:A:650:ARG:H	1.69	0.56
1:A:520:GLU:OE2	1:A:630:ASP:HA	2.06	0.56
1:A:648:ASN:HB2	3:A:802:SO4:O1	2.05	0.55
1:A:531:LYS:NZ	4:A:908:HOH:O	2.38	0.55
1:A:512:LYS:NZ	3:A:805:SO4:O2	2.33	0.55
1:A:610:HIS:O	1:A:611:ARG:HB2	2.06	0.54
1:A:516:ASP:OD2	1:A:638:HIS:NE2	2.29	0.54
1:A:683:PHE:CZ	1:A:718:LEU:HD22	2.46	0.51
1:A:453:ASP:OD2	4:A:903:HOH:O	2.20	0.48
1:A:718:LEU:HD11	1:A:746:VAL:HG11	1.96	0.48
1:A:635:ARG:HD2	3:A:803:SO4:O3	2.15	0.47
1:A:640:ILE:HD11	3:A:803:SO4:O4	2.15	0.46
1:A:457:ASP:OD2	1:A:460:TRP:HD1	1.99	0.45
1:A:586:PHE:CE1	1:A:715:PRO:HG2	2.51	0.44
1:A:594:TYR:CE2	1:A:598:ARG:HD2	2.53	0.44
1:A:506:LYS:HB3	1:A:506:LYS:NZ	2.31	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:578:ARG:HG3	1:A:578:ARG:HH11	1.83	0.43
1:A:638:HIS:HB2	4:A:904:HOH:O	2.18	0.43
1:A:644:LYS:HB2	1:A:644:LYS:HE3	1.74	0.43
1:A:718:LEU:O	1:A:722:MET:HG3	2.19	0.42
1:A:646:THR:HG21	3:A:802:SO4:O3	2.21	0.41
1:A:590:VAL:HG12	1:A:747:LEU:HD13	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	290/310 (94%)	285 (98%)	5 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	250/263 (95%)	240 (96%)	10 (4%)	28 45

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	452	LEU
1	A	459	LEU
1	A	480	GLN
1	A	482	VAL
1	A	506	LYS
1	A	560	GLU
1	A	616	ARG
1	A	646	THR
1	A	703	ARG
1	A	750	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	648	ASN
1	A	713	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
3	SO4	A	802	-	4,4,4	0.43	0	6,6,6	0.68	0
3	SO4	A	805	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	A	804	-	4,4,4	0.13	0	6,6,6	0.04	0
3	SO4	A	803	-	4,4,4	0.16	0	6,6,6	0.10	0
2	A1E09	A	801	1	35,35,36	1.51	6 (17%)	44,49,50	1.29	6 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1E09	A	801	1	-	12/20/20/22	0/4/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	A1E09	C30-N29	4.96	1.45	1.36
2	A	801	A1E09	C12-N10	2.78	1.42	1.37
2	A	801	A1E09	C11-C05	2.49	1.55	1.49
2	A	801	A1E09	C32-C30	2.24	1.55	1.50
2	A	801	A1E09	C18-N29	2.02	1.45	1.41
2	A	801	A1E09	C04-C03	2.01	1.42	1.38

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	801	A1E09	C18-N29-C30	-3.58	121.49	127.99
2	A	801	A1E09	C32-C30-N29	3.05	119.45	114.98
2	A	801	A1E09	C03-C02-C09	2.61	136.00	131.44
2	A	801	A1E09	C01-N07-N08	-2.42	109.89	111.92
2	A	801	A1E09	C02-C01-N07	2.20	108.29	106.48
2	A	801	A1E09	C06-C01-C02	-2.19	120.65	122.45

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	A1E09	C20-C21-O27-C28
2	A	801	A1E09	C22-C21-O27-C28
2	A	801	A1E09	C06-C05-C11-C20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	801	A1E09	C04-C05-C11-C20
2	A	801	A1E09	C06-C05-C11-C24
2	A	801	A1E09	C04-C05-C11-C24
2	A	801	A1E09	C17-C18-N29-C30
2	A	801	A1E09	C19-C18-N29-C30
2	A	801	A1E09	O14-C12-C13-C19
2	A	801	A1E09	N10-C12-C13-C19
2	A	801	A1E09	O14-C12-C13-C15
2	A	801	A1E09	N10-C12-C13-C15


There are no ring outliers.

3 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	802	SO4	3	0
3	A	805	SO4	1	0
3	A	803	SO4	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand A1E09 A 801




Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	294/310 (94%)	0.39	16 (5%)	31	28	20, 33, 56, 69	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	568	PRO	3.3
1	A	451	SER	3.3
1	A	452	LEU	3.2
1	A	577	PRO	2.8
1	A	492	ALA	2.8
1	A	696	GLU	2.7
1	A	580	SER	2.7
1	A	703	ARG	2.5
1	A	453	ASP	2.3
1	A	662	PHE	2.2
1	A	752	GLU	2.1
1	A	470	GLY	2.1
1	A	498	SER	2.1
1	A	454	LEU	2.0
1	A	581	GLU	2.0
1	A	704	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	A1E09	A	801	32/33	0.85	0.12	29,35,51,54	0
3	SO4	A	804	5/5	0.86	0.09	40,43,52,64	0
3	SO4	A	802	5/5	0.87	0.14	50,52,65,66	0
3	SO4	A	803	5/5	0.93	0.08	52,52,59,59	0
3	SO4	A	805	5/5	0.93	0.07	56,58,64,70	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1E09 A 801:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.