Full wwPDB X-ray Structure Validation Report (i)

Nov 10, 2026 – 12:08 PM EST

PDB ID	:	9YW3 / pdb_00009yw3
Title	:	human Siderocalin bound to 343HOPO-Yb
Authors	:	Rupert, P.B.; Strong, R.K.
Deposited on	:	2025-10-23
Resolution	:	2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	178	4% 94% 6%
1	B	178	79% 92% • 5%
1	C	178	2% 91% 6% •

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 4319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neutrophil gelatinase-associated lipocalin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	178	Total	C	N	O	S	0	0	0
			1421	920	235	262	4			
1	B	169	Total	C	N	O	S	0	0	0
			1308	856	212	236	4			
1	C	173	Total	C	N	O	S	0	1	0
			1374	893	226	251	4			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	87	SER	CYS	conflict	UNP P80188
B	87	SER	CYS	conflict	UNP P80188
C	87	SER	CYS	conflict	UNP P80188

- Molecule 2 is YTTERBIUM (III) ION (CCD ID: YB) (formula: Yb).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Yb	0	0
			1	1		
2	B	1	Total	Yb	0	0
			1	1		
2	C	1	Total	Yb	0	0
			1	1		

- Molecule 3 is N,N'-butane-1,4-diylbis[1-hydroxy-N-(3-{[(1-hydroxy-6-oxo-1,6-dihydropyridin-2-yl)carbonyl]amino}propyl)-6-oxo-1,6-dihydropyridine-2-carboxamide] (CCD ID: 4OL) (formula: $C_{34}H_{38}N_8O_{12}$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 54 34 8 12	0	0
3	B	1	Total C N O 54 34 8 12	0	0
3	C	1	Total C N O 54 34 8 12	0	0

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0
4	C	1	Total Cl 1 1	0	0

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	C	1	Total O S 5 4 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	13	Total O 13 13	0	0
6	B	1	Total O 1 1	0	0
6	C	25	Total O 25 25	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Neutrophil gelatinase-associated lipocalin


Chain A:








• Molecule 1: Neutrophil gelatinase-associated lipocalin

Chain B:









• Molecule 1: Neutrophil gelatinase-associated lipocalin

Chain C:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α, β, γ	115.25Å 115.25Å 118.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.57 – 2.55 38.57 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.9 (38.57-2.55) 99.9 (38.57-2.55)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.32 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.218 , 0.240 0.219 , 0.237	Depositor DCC
R_{free} test set	1271 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å^2)	49.1	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 42.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.021 for -h,l,k 0.015 for -l,-k,-h	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	4319	wwPDB-VP
Average B, all atoms (Å^2)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 4OL, YB, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.49	0/1458	0.80	0/1979
1	B	0.48	0/1344	0.76	0/1834
1	C	0.49	0/1414	0.79	0/1925
All	All	0.49	0/4216	0.78	0/5738

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1421	0	1394	6	0
1	B	1308	0	1243	3	0
1	C	1374	0	1330	5	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	54	0	34	2	0
3	B	54	0	35	2	0
3	C	54	0	34	0	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	1	0	0	0	0
5	A	5	0	0	0	0
5	C	5	0	0	1	0
6	A	13	0	0	0	0
6	B	1	0	0	1	0
6	C	25	0	0	0	0
All	All	4319	0	4070	16	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:301:4OL:O9	6:B:401:HOH:O	1.98	0.81
1:C:25:ASN:ND2	5:C:204:SO4:O4	2.32	0.60
1:A:100:TYR:CD2	3:A:202:4OL:H32	2.43	0.53
1:A:91:GLU:OE1	1:A:109:ARG:NH1	2.42	0.53
1:A:33:VAL:HG21	1:A:52:TYR:CE2	2.47	0.49
1:C:33:VAL:HG21	1:C:52:TYR:CE2	2.47	0.49
1:A:41:ILE:HD11	3:A:202:4OL:H4	1.94	0.49
1:B:33:VAL:HG21	1:B:52:TYR:CE2	2.48	0.48
1:C:91:GLU:OE1	1:C:109:ARG:NH2	2.46	0.47
1:C:69:VAL:HG22	1:C:78:TYR:CD1	2.52	0.45
1:A:69:VAL:HG22	1:A:78:TYR:CD1	2.52	0.44
1:B:69:VAL:HG22	1:B:78:TYR:CD1	2.52	0.44
1:B:31:TRP:CE3	1:B:138:TYR:HB3	2.54	0.42
1:A:31:TRP:CE3	1:A:138:TYR:HB3	2.54	0.42
3:B:301:4OL:C22	3:B:301:4OL:H6	2.50	0.41
1:C:31:TRP:CE3	1:C:138:TYR:HB3	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:150:GLU:OE1	1:C:150:GLU:OE1[8_665]	2.04	0.16

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	176/178 (99%)	173 (98%)	3 (2%)	0	100	100
1	B	165/178 (93%)	162 (98%)	3 (2%)	0	100	100
1	C	172/178 (97%)	169 (98%)	3 (2%)	0	100	100
All	All	513/534 (96%)	504 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/163 (96%)	156 (100%)	0	100	100
1	B	137/163 (84%)	137 (100%)	0	100	100
1	C	149/163 (91%)	148 (99%)	1 (1%)	76	85
All	All	442/489 (90%)	441 (100%)	1 (0%)	87	93

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	163	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	ASN
1	A	117	GLN
1	B	25	ASN
1	B	49	GLN
1	C	128	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	4OL	A	202	2	57,57,57	2.73	11 (19%)	64,78,78	1.79	11 (17%)
3	4OL	B	301	2	57,57,57	2.59	11 (19%)	64,78,78	1.55	7 (10%)
5	SO4	A	204	-	4,4,4	0.32	0	6,6,6	0.08	0
5	SO4	C	204	-	4,4,4	0.33	0	6,6,6	0.08	0
3	4OL	C	202	2	57,57,57	1.87	10 (17%)	64,78,78	1.24	6 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	4OL	C	202	2	-	3/45/45/45	0/4/4/4
3	4OL	A	202	2	-	11/45/45/45	0/4/4/4
3	4OL	B	301	2	-	10/45/45/45	0/4/4/4

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	202	4OL	C44-N45	9.79	1.42	1.39
3	A	202	4OL	C26-N27	8.97	1.42	1.39
3	B	301	4OL	C44-N45	8.46	1.42	1.39
3	B	301	4OL	C4-N3	8.26	1.41	1.39
3	A	202	4OL	C36-N35	8.03	1.41	1.39
3	B	301	4OL	C36-N35	8.01	1.41	1.39
3	B	301	4OL	C26-N27	6.86	1.41	1.39
3	A	202	4OL	C4-N3	6.15	1.41	1.39
3	A	202	4OL	O48-N27	5.47	1.44	1.38
3	C	202	4OL	O50-N45	5.31	1.44	1.38
3	A	202	4OL	O10-N3	4.83	1.44	1.38
3	A	202	4OL	O46-N35	4.77	1.44	1.38
3	C	202	4OL	C4-N3	4.74	1.40	1.39
3	A	202	4OL	O50-N45	4.65	1.44	1.38
3	C	202	4OL	O10-N3	4.61	1.43	1.38
3	B	301	4OL	O48-N27	4.60	1.43	1.38
3	B	301	4OL	O50-N45	4.51	1.43	1.38
3	B	301	4OL	O46-N35	4.50	1.43	1.38
3	B	301	4OL	O10-N3	4.50	1.43	1.38
3	C	202	4OL	C36-N35	4.49	1.40	1.39
3	C	202	4OL	O48-N27	4.34	1.43	1.38
3	C	202	4OL	O46-N35	4.08	1.43	1.38
3	C	202	4OL	C44-N45	3.93	1.40	1.39
3	C	202	4OL	C26-N27	3.37	1.40	1.39
3	A	202	4OL	C22-N27	2.65	1.42	1.38
3	A	202	4OL	C40-N45	2.34	1.42	1.38
3	B	301	4OL	C2-N3	2.33	1.42	1.38
3	B	301	4OL	C22-N27	2.31	1.42	1.38
3	B	301	4OL	C40-N45	2.18	1.41	1.38
3	C	202	4OL	C40-N45	2.17	1.41	1.38
3	C	202	4OL	C2-N3	2.13	1.41	1.38
3	A	202	4OL	C34-N35	2.07	1.41	1.38

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	202	4OL	O10-N3-C4	5.86	120.95	116.75
3	A	202	4OL	O50-N45-C44	5.44	120.65	116.75
3	A	202	4OL	O46-N35-C36	5.41	120.63	116.75
3	B	301	4OL	O46-N35-C36	5.20	120.48	116.75
3	B	301	4OL	O50-N45-C44	4.68	120.11	116.75
3	A	202	4OL	O48-N27-C26	4.65	120.09	116.75
3	C	202	4OL	C22-C16-N15	3.91	124.31	118.45
3	B	301	4OL	O10-N3-C4	3.86	119.52	116.75
3	B	301	4OL	O48-N27-C26	3.65	119.37	116.75
3	C	202	4OL	O48-N27-C26	3.34	119.15	116.75
3	B	301	4OL	C34-C28-N21	3.21	123.26	118.45
3	A	202	4OL	C22-C16-N15	2.96	122.88	118.45
3	A	202	4OL	C34-C28-N21	2.92	122.83	118.45
3	A	202	4OL	O51-C44-N45	2.67	121.38	118.76
3	B	301	4OL	C22-C16-N15	2.63	122.39	118.45
3	C	202	4OL	C41-C40-N45	2.59	120.43	118.48
3	C	202	4OL	O46-N35-C36	2.53	118.56	116.75
3	C	202	4OL	O10-N3-C4	2.42	118.49	116.75
3	A	202	4OL	C7-C2-N3	2.20	120.14	118.48
3	C	202	4OL	C14-N15-C16	2.15	126.61	120.66
3	A	202	4OL	O51-C44-C43	-2.09	118.42	124.35
3	A	202	4OL	O47-C36-N35	2.08	120.80	118.76
3	B	301	4OL	O51-C44-N45	2.08	120.80	118.76
3	A	202	4OL	C1-C2-N3	-2.02	118.35	122.27

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	301	4OL	O53-C33-C40-N45
3	B	301	4OL	O53-C33-C40-C41
3	A	202	4OL	C18-C19-C20-N21
3	C	202	4OL	N15-C17-C18-C19
3	B	301	4OL	N32-C33-C40-N45
3	B	301	4OL	N32-C33-C40-C41
3	A	202	4OL	O54-C16-N15-C17
3	B	301	4OL	O54-C16-N15-C17
3	A	202	4OL	C22-C16-N15-C17
3	B	301	4OL	C22-C16-N15-C17
3	B	301	4OL	C18-C19-C20-N21
3	A	202	4OL	N9-C1-C2-N3
3	A	202	4OL	O8-C1-C2-N3
3	A	202	4OL	O53-C33-C40-N45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	202	4OL	O8-C1-C2-C7
3	A	202	4OL	O53-C33-C40-C41
3	A	202	4OL	N9-C1-C2-C7
3	A	202	4OL	N32-C33-C40-N45
3	C	202	4OL	O54-C16-C22-N27
3	B	301	4OL	O54-C16-C22-C23
3	A	202	4OL	O55-C28-C34-N35
3	B	301	4OL	O55-C28-C34-N35
3	B	301	4OL	O55-C28-C34-C39
3	C	202	4OL	O54-C16-C22-C23


There are no ring outliers.

3 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	202	4OL	2	0
3	B	301	4OL	2	0
5	C	204	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand 4OL A 202




Bond lengths



Bond angles




Torsions




Rings


Ligand 4OL B 301




Bond lengths



Bond angles




Torsions



Rings

Ligand 4OL C 202



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	178/178 (100%)	0.36	8 (4%) 38 38	31, 49, 72, 109	0
1	B	169/178 (94%)	3.09	140 (82%) 0 0	99, 129, 155, 170	0
1	C	173/178 (97%)	0.02	3 (1%) 69 69	21, 36, 59, 105	1 (0%)
All	All	520/534 (97%)	1.13	151 (29%) 1 1	21, 53, 146, 170	1 (0%)

All (151) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	5	SER	7.8
1	B	153	ILE	6.9
1	A	178	GLY	6.6
1	B	27	PHE	6.0
1	B	150	GLU	5.8
1	B	60	GLU	5.7
1	B	6	ASP	5.7
1	B	42	LEU	5.6
1	B	152	PHE	5.5
1	B	64	TYR	5.4
1	B	167	VAL	5.2
1	B	89	PRO	5.0
1	B	20	GLN	5.0
1	B	84	VAL	4.7
1	B	85	PRO	4.7
1	B	48	PRO	4.6
1	B	144	LEU	4.6
1	B	148	LEU	4.6
1	B	47	ASP	4.5
1	B	113	THR	4.4
1	B	22	PHE	4.4
1	B	115	TYR	4.4
1	B	62	LYS	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	154	ARG	4.3
1	B	92	PHE	4.3
1	B	23	GLN	4.2
1	B	135	ILE	4.2
1	B	136	THR	4.2
1	B	110	VAL	4.2
1	B	97	ILE	4.2
1	B	116	ASN	4.1
1	B	34	VAL	4.1
1	B	18	LEU	4.0
1	B	53	ALA	4.0
1	B	32	TYR	4.0
1	B	126	VAL	3.9
1	B	107	LEU	3.9
1	B	41	ILE	3.9
1	B	21	ASN	3.8
1	B	58	LEU	3.8
1	B	30	LYS	3.8
1	B	13	LEU	3.8
1	B	19	GLN	3.8
1	B	16	VAL	3.7
1	B	59	LYS	3.7
1	B	94	LEU	3.7
1	B	25	ASN	3.7
1	B	170	VAL	3.6
1	B	12	PRO	3.6
1	B	176	ILE	3.6
1	B	175	CYS	3.6
1	B	172	ILE	3.6
1	B	143	GLU	3.5
1	B	17	PRO	3.5
1	B	11	PRO	3.5
1	B	141	THR	3.4
1	B	119	ALA	3.4
1	B	70	LEU	3.4
1	B	137	LEU	3.4
1	B	40	ALA	3.4
1	B	49	GLN	3.3
1	B	63	SER	3.3
1	B	15	LYS	3.3
1	B	39	ASN	3.3
1	B	87	SER	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	66	VAL	3.3
1	B	149	LYS	3.3
1	B	138	TYR	3.2
1	B	93	THR	3.2
1	B	145	THR	3.2
1	B	158	SER	3.2
1	B	168	PHE	3.2
1	B	65	ASN	3.2
1	B	80	ILE	3.2
1	B	174	GLN	3.1
1	B	161	LEU	3.1
1	B	111	VAL	3.1
1	B	103	LEU	3.1
1	B	156	SER	3.1
1	B	160	GLY	3.1
1	B	104	THR	3.0
1	B	164	ASN	3.0
1	B	142	LYS	3.0
1	B	90	GLY	3.0
1	B	118	HIS	3.0
1	B	122	PHE	3.0
1	B	129	ASN	3.0
1	B	50	LYS	2.9
1	B	78	TYR	2.9
1	B	96	ASN	2.9
1	B	131	GLU	2.9
1	B	7	LEU	2.9
1	B	166	ILE	2.9
1	A	1	GLN	2.9
1	B	95	GLY	2.9
1	B	109	ARG	2.8
1	B	162	PRO	2.8
1	B	171	PRO	2.8
1	A	150	GLU	2.8
1	B	121	VAL	2.8
1	B	67	THR	2.8
1	B	159	LEU	2.8
1	B	155	PHE	2.8
1	B	79	TRP	2.8
1	B	120	MET	2.8
1	A	80	ILE	2.7
1	B	8	ILE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	144	LEU	2.7
1	B	35	GLY	2.7
1	B	112	SER	2.7
1	B	88	GLN	2.7
1	B	177	ASP	2.7
1	B	33	VAL	2.7
1	B	36	LEU	2.7
1	B	147	GLU	2.7
1	B	82	THR	2.6
1	B	52	TYR	2.6
1	B	134	LYS	2.6
1	B	56	TYR	2.6
1	B	9	PRO	2.6
1	B	151	ASN	2.6
1	B	83	PHE	2.5
1	B	132	TYR	2.5
1	B	77	ASP	2.5
1	B	24	ASP	2.5
1	B	71	PHE	2.4
1	A	79	TRP	2.4
1	B	26	GLN	2.4
1	B	14	SER	2.4
1	B	69	VAL	2.4
1	B	55	ILE	2.3
1	B	169	PRO	2.3
1	B	31	TRP	2.3
1	B	133	PHE	2.3
1	A	60	GLU	2.3
1	B	114	ASN	2.3
1	B	86	GLY	2.2
1	B	68	SER	2.2
1	B	76	CYS	2.2
1	A	46	LYS	2.2
1	B	101	PRO	2.2
1	B	128	GLN	2.1
1	C	44	GLU	2.1
1	B	139	GLY	2.1
1	B	61	ASP	2.1
1	B	105	SER	2.1
1	A	43	ARG	2.1
1	C	97	ILE	2.1
1	B	75	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	100	TYR	2.0
1	B	123	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	SO4	C	204	5/5	0.66	0.15	108,110,113,114	0
5	SO4	A	204	5/5	0.80	0.22	85,86,87,87	0
3	4OL	B	301	54/54	0.84	0.15	108,127,137,139	0
3	4OL	A	202	54/54	0.87	0.15	53,80,90,95	0
3	4OL	C	202	54/54	0.97	0.08	38,46,54,56	0
4	CL	C	203	1/1	0.98	0.08	36,36,36,36	0
2	YB	B	300	1/1	0.99	0.03	127,127,127,127	0
2	YB	A	201	1/1	0.99	0.07	77,77,77,77	0
4	CL	A	203	1/1	0.99	0.04	49,49,49,49	0
2	YB	C	201	1/1	1.00	0.02	46,46,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around 4OL B 301:


$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around 4OL A 202:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)








Electron density around 4OL C 202:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.