



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 16, 2026 – 06:12 PM EDT

PDB ID : 9YXY / pdb\_00009yxy  
Title : Crystal structure of EEPD1 EEP domain dimer at pH 5.5  
Authors : Shen, R.; Arvai, A.S.; Brosey, C.A.; Tsai, C.; Tsutakawa, S.E.; Tainer, J.A.  
Deposited on : 2025-10-27  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.48.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	 4% (Poor fit), 82% (0-1 outliers), 7% (2 outliers), 11% (3+ outliers or not modelled)
1	B	302	 2% (Poor fit), 78% (0-1 outliers), 8% (2 outliers), 15% (3+ outliers or not modelled)

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8446 atoms, of which 4107 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endonuclease/exonuclease/phosphatase family domain-containing protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	268	Total	C	H	N	O	S	0	4	0
			4156	1333	2065	367	384	7			
1	B	258	Total	C	H	N	O	S	0	4	0
			4066	1302	2023	359	375	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	242	GLY	-	expression tag	UNP Q7L9B9
A	243	PRO	-	expression tag	UNP Q7L9B9
A	244	MET	-	expression tag	UNP Q7L9B9
A	343	ASN	SER	variant	UNP Q7L9B9
B	242	GLY	-	expression tag	UNP Q7L9B9
B	243	PRO	-	expression tag	UNP Q7L9B9
B	244	MET	-	expression tag	UNP Q7L9B9
B	343	ASN	SER	variant	UNP Q7L9B9

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	14	3	8	3	0	0

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	10	2	6	2	0	0

- Molecule 4 is IMIDAZOLE (CCD ID: IMD) (formula:  $C_3H_5N_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	N		
4	B	1	10	3	5	2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	107	107	107	0	0
5	B	81	83	83	0	2


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endonuclease/exonuclease/phosphatase family domain-containing protein 1



- Molecule 1: Endonuclease/exonuclease/phosphatase family domain-containing protein 1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	38.94Å 116.32Å 121.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.24 – 2.00 38.24 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.7 (38.24-2.00) 85.5 (38.24-2.00)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.87 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, $R_{free}$	0.194 , 0.232 0.194 , 0.232	Depositor DCC
$R_{free}$ test set	1805 reflections (3.46%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.4	Xtrriage
Anisotropy	0.292	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 54.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.013 for -h,l,k	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8446	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, GOL, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.15	0/2169	0.38	0/2943
1	B	0.16	0/2110	0.38	0/2863
All	All	0.16	0/4279	0.38	0/5806

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2091	2065	2040	14	0
1	B	2043	2023	2003	21	0
2	A	6	8	8	0	0
3	A	4	6	6	0	0
4	B	5	5	5	0	0
5	A	107	0	0	0	0
5	B	83	0	0	0	0
All	All	4339	4107	4062	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:393:VAL:HG21	1:B:398:LEU:HD12	1.50	0.92
1:B:362:MET:HE1	1:B:393:VAL:HG22	1.61	0.82
1:B:340:LYS:HD3	1:B:369:SER:HB2	1.77	0.66
1:B:393:VAL:CG2	1:B:398:LEU:HD12	2.25	0.65
1:B:261:LEU:HD11	1:B:362:MET:SD	2.38	0.63
1:B:362:MET:SD	1:B:393:VAL:HG13	2.39	0.62
1:B:255:ARG:HD3	1:B:503:THR:HB	1.83	0.60
1:B:261:LEU:C	1:B:261:LEU:HD23	2.28	0.59
1:B:296:LEU:HD22	1:B:362:MET:CE	2.33	0.59
1:B:362:MET:CE	1:B:393:VAL:HG22	2.33	0.58
1:B:459:ASP:O	1:B:463:LYS:HG3	2.09	0.52
1:A:363:GLU:CG	1:A:392:LYS:HB3	2.42	0.49
1:B:296:LEU:HD22	1:B:362:MET:HE1	1.93	0.49
1:A:344:GLN:NE2	1:A:383:GLY:N	2.61	0.48
1:B:340:LYS:NZ	1:B:371:GLU:HG3	2.29	0.47
1:A:370:GLN:HG2	1:A:388:LEU:HD13	1.96	0.47
1:A:344:GLN:HG2	1:A:345:LEU:N	2.30	0.46
1:B:370:GLN:HG3	1:B:388:LEU:HD13	1.98	0.46
1:B:370:GLN:HG3	1:B:436:THR:HG22	1.98	0.46
1:A:390:ARG:NH1	1:A:440:GLU:OE2	2.50	0.44
1:B:370:GLN:CG	1:B:388:LEU:HD13	2.48	0.44
1:B:535:LEU:HD12	1:B:535:LEU:C	2.43	0.43
1:B:261:LEU:CD1	1:B:362:MET:SD	3.06	0.43
1:A:432:THR:O	1:A:436:THR:HG23	2.18	0.43
1:A:370:GLN:CG	1:A:388:LEU:HD13	2.49	0.42
1:B:295:LYS:HD2	1:B:360:ALA:HB2	2.02	0.42
1:A:468:HIS:HB3	1:A:493:TRP:CE2	2.55	0.42
1:A:366:ASP:OD1	1:A:366:ASP:C	2.62	0.41
1:A:446:LEU:HD23	1:A:446:LEU:H	1.86	0.41
1:A:388:LEU:HD12	1:A:388:LEU:N	2.35	0.41
1:A:261:LEU:HD11	1:A:362:MET:SD	2.61	0.41
1:B:502:PHE:CE2	1:B:504:GLY:HA2	2.56	0.40
1:A:338:ALA:HB2	1:A:387:TYR:CD1	2.55	0.40
1:B:429:PHE:CZ	1:B:433:LEU:HG	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/302 (88%)	258 (97%)	7 (3%)	1 (0%)	30	27
1	B	256/302 (85%)	246 (96%)	9 (4%)	1 (0%)	30	27
All	All	522/604 (86%)	504 (97%)	16 (3%)	2 (0%)	30	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	269	GLN
1	A	269	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/251 (89%)	223 (100%)	0	100	100
1	B	220/251 (88%)	220 (100%)	0	100	100
All	All	443/502 (88%)	443 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	473	HIS
1	B	343	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	434	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	IMD	B	601	-	3,5,5	0.16	0	4,5,5	1.00	0
3	EDO	A	602	-	3,3,3	0.28	0	2,2,2	0.30	0
2	GOL	A	601	-	5,5,5	0.39	0	5,5,5	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IMD	B	601	-	-	-	0/1/1/1
3	EDO	A	602	-	-	0/1/1/1	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	601	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	GOL	C1-C2-C3-O3
2	A	601	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	268/302 (88%)	0.09	11 (4%) 42 40	27, 53, 107, 128	2 (0%)
1	B	258/302 (85%)	0.07	6 (2%) 61 59	26, 56, 111, 144	2 (0%)
All	All	526/604 (87%)	0.08	17 (3%) 50 48	26, 54, 109, 144	4 (0%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	429	PHE	4.4
1	A	541	GLU	3.8
1	A	371	GLU	3.6
1	B	408	LEU	3.3
1	A	398	LEU	3.0
1	A	437	LEU	2.8
1	A	433	LEU	2.7
1	A	425	ARG	2.7
1	B	254	THR	2.4
1	A	407	ALA	2.3
1	B	255	ARG	2.2
1	A	430	ALA	2.1
1	B	437	LEU	2.1
1	B	429	PHE	2.1
1	B	432	THR	2.1
1	A	436	THR	2.0
1	A	406	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	A	602	4/4	0.79	0.15	51,63,71,76	0
4	IMD	B	601	5/5	0.88	0.13	54,64,73,76	0
2	GOL	A	601	6/6	0.95	0.07	40,51,64,77	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.