



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2026 – 02:06 PM EST

PDB ID : 9Z44 / pdb_00009z44
Title : Isoreticular co-crystal 1 with symmetrical expanded duplex (31mer) containing insert sequence CCCGGCCGGA and loaded with C-clamp domain
Authors : Shields, E.T.; Slaughter, C.K.; Magna, E.N.; Snow, C.D.
Deposited on : 2025-11-08
Resolution : 7.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 7.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1107 (10.00-4.00)
Clashscore	180529	1146 (10.00-4.00)
Ramachandran outliers	177936	1014 (10.00-4.00)
Sidechain outliers	177891	1035 (10.00-3.96)
RSRZ outliers	164620	1102 (10.00-4.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	31	 61% 39%
2	B	31	 68% 32%
3	C	263	 10% 78% 5% 17%
4	D	33	 55% 42%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2821 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (31-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	31	Total	C	N	O	P	0	0	0
			630	298	116	185	31			

- Molecule 2 is a DNA chain called DNA (31-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	31	Total	C	N	O	P	0	0	0
			641	302	121	187	31			

- Molecule 3 is a protein called Replication initiation protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	218	Total	C	N	O	S	0	0	0
			1438	886	259	290	3			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-11	MET	-	initiating methionine	UNP P03856
C	-10	ARG	-	expression tag	UNP P03856
C	-9	GLY	-	expression tag	UNP P03856
C	-8	SER	-	expression tag	UNP P03856
C	-7	HIS	-	expression tag	UNP P03856
C	-6	HIS	-	expression tag	UNP P03856
C	-5	HIS	-	expression tag	UNP P03856
C	-4	HIS	-	expression tag	UNP P03856
C	-3	HIS	-	expression tag	UNP P03856
C	-2	HIS	-	expression tag	UNP P03856
C	-1	GLY	-	expression tag	UNP P03856
C	0	SER	-	expression tag	UNP P03856
C	118	PRO	ARG	conflict	UNP P03856

- Molecule 4 is a protein called SLC2A4 regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	19	Total	C	N	O	S	0	0	0
			112	67	22	20	3			

- Molecule 1: DNA (31-MER)



- | | | | |
|-----|------------|--------------------------|--------------------------|
| G17 | G27
G28 | G36
T37
C38
A39 | T44
C45
C46
G47 |
|-----|------------|--------------------------|--------------------------|

- | |
|-----|
| MET | ARG | GLY | SER | HIS | HIS | HIS | HIS | HIS | GLY | SER | MET | ALA | GLU | THR | ALA | VAL | ILE | ASN | LYS | ARG | K13 | M14 | S15 | S21 | A28 | Y29 | S30 | L31 | S32 | Q35 | K36 | R37 | M38 | L39 | Y40 | F42 | F43 | S49 | ASP | GLY | THR | LEU | GLN | GLU | H56 | H63 | Y67 | T70 |
|-----|



- GLY
ASP
ALA
LYS
LYS
C360
W378
LYS
LYS
ALA
CYS
GLN
ARG
PHE
LEU
ASP

4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	73.49Å 127.65Å 141.18Å 90.00° 91.98° 90.00°	Depositor
Resolution (Å)	47.99 – 7.20 47.99 – 7.20	Depositor EDS
% Data completeness (in resolution range)	98.1 (47.99-7.20) 97.8 (47.99-7.20)	Depositor EDS
R_{merge}	2.02	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 7.37Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.280 , 0.345 0.280 , 0.343	Depositor DCC
R_{free} test set	192 reflections (9.91%)	wwPDB-VP
Wilson B-factor (Å ²)	321.8	Xtriage
Anisotropy	0.258	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.23 , 299.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	0.129 for h,-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2821	wwPDB-VP
Average B, all atoms (Å ²)	309.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.19	0/705	0.37	0/1084
2	B	0.18	0/719	0.40	0/1109
3	C	0.14	0/1462	0.46	0/2010
4	D	0.10	0/112	0.27	0/151
All	All	0.16	0/2998	0.42	0/4354

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	630	0	347	9	0
2	B	641	0	348	6	0
3	C	1438	0	1123	6	0
4	D	112	0	72	1	0
All	All	2821	0	1890	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:DG:H3'	4:D:378:TRP:HA	1.67	0.76
2:B:27:DG:H2'	2:B:28:DG:C8	2.33	0.62
3:C:28:ALA:HA	3:C:191:ARG:O	2.08	0.54
1:A:-3:DC:H2''	1:A:-2:DG:C8	2.44	0.52
3:C:15:SER:OG	3:C:151:LEU:HB3	2.09	0.52
1:A:-3:DC:H2''	1:A:-2:DG:H5'	1.92	0.52
1:A:3:DT:H2'	1:A:4:DG:C8	2.49	0.47
3:C:180:ILE:HA	3:C:240:PHE:O	2.14	0.47
1:A:23:DC:H2''	1:A:24:DC:C5	2.49	0.47
2:B:36:DG:H1'	2:B:37:DT:H5'	1.96	0.47
3:C:117:LYS:HB2	3:C:130:HIS:HB2	1.96	0.46
1:A:0:DA:H2''	1:A:1:DC:C6	2.52	0.44
3:C:98:ARG:N	3:C:99:PRO:HD3	2.33	0.43
2:B:46:DC:H2''	2:B:47:DG:C8	2.54	0.42
3:C:21:SER:HB2	3:C:141:GLY:HA3	2.00	0.42
1:A:-4:DC:H6	1:A:-4:DC:H2'	1.71	0.42
2:B:38:DC:H2''	2:B:39:DA:H8	1.85	0.42
2:B:44:DT:H2''	2:B:45:DC:C5	2.55	0.41
2:B:45:DC:H2''	2:B:46:DC:C5	2.55	0.41
1:A:-1:DG:H2''	1:A:0:DA:C8	2.56	0.41
1:A:20:DA:H2''	1:A:21:DG:H8	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	210/263 (80%)	203 (97%)	5 (2%)	2 (1%)	13	49
4	D	17/33 (52%)	15 (88%)	2 (12%)	0	100	100
All	All	227/296 (77%)	218 (96%)	7 (3%)	2 (1%)	14	52

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	99	PRO
3	C	227	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	115/236 (49%)	115 (100%)	0	100	100
4	D	5/28 (18%)	5 (100%)	0	100	100
All	All	120/264 (46%)	120 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	31/31 (100%)	-0.20	0	100 100	210, 321, 391, 394	0
2	B	31/31 (100%)	-0.19	0	100 100	232, 309, 368, 382	0
3	C	218/263 (82%)	0.44	27 (12%)	9 14	196, 290, 420, 611	0
4	D	19/33 (57%)	-0.60	0	100 100	360, 385, 462, 555	0
All	All	299/358 (83%)	0.25	27 (9%)	17 19	196, 297, 434, 611	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	39	LEU	5.6
3	C	86	LEU	4.4
3	C	109	TYR	4.0
3	C	160	PRO	3.4
3	C	31	LEU	3.4
3	C	123	SER	3.3
3	C	38	MET	3.1
3	C	49	SER	2.9
3	C	129	VAL	2.8
3	C	40	TYR	2.8
3	C	159	ASN	2.6
3	C	30	SER	2.6
3	C	32	SER	2.6
3	C	128	SER	2.5
3	C	67	TYR	2.4
3	C	63	HIS	2.4
3	C	82	ILE	2.3
3	C	88	SER	2.3
3	C	43	VAL	2.3
3	C	35	GLN	2.2
3	C	72	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	213	VAL	2.2
3	C	42	PHE	2.1
3	C	186	ASP	2.1
3	C	37	ARG	2.1
3	C	89	PHE	2.1
3	C	87	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.