Full wwPDB X-ray Structure Validation Report (i)

Nov 1, 2026 – 12:05 PM EST

PDB ID	:	9Z59 / pdb_00009z59
Title	:	Crystal structure of MPXV-A35 in complex with antibody A35-Ab10
Authors	:	Zhang, D.; Wilson, I.A.
Deposited on	:	2025-11-11
Resolution	:	2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION





The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	228	
2	L	216	
3	C	100	
3	D	100	

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 4558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called A35-Ab10 antibody heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	228	Total	C	N	O	S	0	0	0
			1686	1063	288	329	6			

- Molecule 2 is a protein called A35-Ab10 antibody light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	208	Total	C	N	O	S	0	0	0
			1601	1000	278	317	6			

- Molecule 3 is a protein called Protein OPG161.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	75	Total	C	N	O	S	0	0	0
			593	372	93	124	4			
3	D	73	Total	C	N	O	S	0	0	0
			572	358	90	120	4			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	182	GLY	-	expression tag	UNP A0A7H0DND2
C	183	SER	-	expression tag	UNP A0A7H0DND2
C	184	HIS	-	expression tag	UNP A0A7H0DND2
C	185	HIS	-	expression tag	UNP A0A7H0DND2
C	186	HIS	-	expression tag	UNP A0A7H0DND2
C	187	HIS	-	expression tag	UNP A0A7H0DND2
C	188	HIS	-	expression tag	UNP A0A7H0DND2
C	189	HIS	-	expression tag	UNP A0A7H0DND2
D	182	GLY	-	expression tag	UNP A0A7H0DND2
D	183	SER	-	expression tag	UNP A0A7H0DND2
D	184	HIS	-	expression tag	UNP A0A7H0DND2
D	185	HIS	-	expression tag	UNP A0A7H0DND2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	186	HIS	-	expression tag	UNP A0A7H0DND2
D	187	HIS	-	expression tag	UNP A0A7H0DND2
D	188	HIS	-	expression tag	UNP A0A7H0DND2
D	189	HIS	-	expression tag	UNP A0A7H0DND2

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	H	1	Total	C	O	0	0
			4	2	2		
4	H	1	Total	C	O	0	0
			4	2	2		
4	H	1	Total	C	O	0	0
			4	2	2		
4	H	1	Total	C	O	0	0
			4	2	2		
4	L	1	Total	C	O	0	0
			4	2	2		
4	L	1	Total	C	O	0	0
			4	2	2		
4	L	1	Total	C	O	0	0
			4	2	2		
4	L	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	H	31	Total O 31 31	0	0
5	L	33	Total O 33 33	0	0
5	C	4	Total O 4 4	0	0
5	D	2	Total O 2 2	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: A35-Ab10 antibody heavy chain

Chain H:




• Molecule 2: A35-Ab10 antibody light chain

Chain L:




• Molecule 3: Protein OPG161

Chain C:



• Molecule 3: Protein OPG161

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α, β, γ	119.53Å 119.53Å 92.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.07 – 2.65 42.07 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.8 (42.07-2.65) 98.0 (42.07-2.65)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.31 (at 2.65Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.179 , 0.231 0.179 , 0.230	Depositor DCC
R_{free} test set	1995 reflections (9.21%)	wwPDB-VP
Wilson B-factor (Å^2)	38.8	Xtriage
Anisotropy	0.074	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 49.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.045 for h,-h-k,-l	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	4558	wwPDB-VP
Average B, all atoms (Å^2)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	H	0.12	0/1730	0.39	0/2363
2	L	0.20	0/1632	0.40	0/2207
3	C	0.09	0/606	0.29	0/823
3	D	0.12	0/585	0.35	0/794
All	All	0.15	0/4553	0.38	0/6187

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	L	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	L	108	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1686	0	1665	14	0
2	L	1601	0	1570	23	0
3	C	593	0	542	6	0
3	D	572	0	517	10	0
4	D	4	0	6	1	0
4	H	16	0	24	0	0
4	L	16	0	24	0	0
5	C	4	0	0	0	0
5	D	2	0	0	0	0
5	H	31	0	0	1	0
5	L	33	0	0	1	0
All	All	4558	0	4348	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:27:VAL:HG21	2:L:92:GLY:HA3	1.79	0.64
1:H:119:PRO:HB3	1:H:145:TYR:HB3	1.83	0.60
1:H:129:LYS:NZ	5:H:401:HOH:O	2.33	0.60
3:C:114:SER:HA	3:C:177:LYS:HD3	1.86	0.58
2:L:35:TRP:HB2	2:L:48:ILE:HB	1.86	0.57
3:D:135:ASN:HD22	3:D:136:LYS:N	2.04	0.55
2:L:27:VAL:HG13	2:L:29:SER:H	1.73	0.52
2:L:155:GLN:OE1	2:L:158:ASN:ND2	2.43	0.52
2:L:6:GLN:NE2	2:L:86:TYR:O	2.39	0.52
3:D:136:LYS:HG3	3:D:140:LEU:HD23	1.91	0.52
2:L:19:ALA:HB2	2:L:78:LEU:HD11	1.91	0.52
1:H:13:LYS:HB2	1:H:16:GLU:HG3	1.92	0.52
2:L:8:PRO:HG3	2:L:11:LEU:HD13	1.91	0.52
3:C:135:ASN:ND2	3:C:137:SER:OG	2.43	0.52
2:L:27:VAL:HB	2:L:95(A):MET:SD	2.50	0.51
3:C:105:TYR:CE2	3:C:106:GLN:HG3	2.47	0.50
3:C:103:LEU:HB2	3:C:110:TYR:HB2	1.92	0.50
3:C:181:THR:O	3:C:181:THR:OG1	2.29	0.50
2:L:163:VAL:HG22	2:L:175:LEU:HD12	1.94	0.49
1:H:11:LEU:HD11	1:H:112:SER:HB3	1.94	0.49
1:H:47:TRP:CD2	2:L:96:LEU:HD23	2.47	0.48
2:L:158:ASN:OD1	2:L:158:ASN:N	2.46	0.48
1:H:9:PRO:HA	1:H:107:THR:HG23	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:29:ILE:HG22	1:H:35(A):TRP:CE2	2.50	0.47
2:L:210:ASN:O	2:L:213:GLU:HG2	2.16	0.46
1:H:116:THR:HG22	1:H:203:SER:HB3	1.98	0.46
1:H:4:LEU:HD22	1:H:24:VAL:HG22	1.97	0.46
1:H:35:HIS:CE1	1:H:95:GLY:HA3	2.51	0.46
1:H:100:MET:SD	3:D:177:LYS:HD2	2.55	0.46
2:L:120:PRO:HD3	2:L:132:VAL:HG22	1.99	0.45
3:C:134:PRO:HD3	3:C:179:PHE:O	2.17	0.45
3:D:135:ASN:O	3:D:139:VAL:HG22	2.17	0.44
3:D:139:VAL:O	3:D:142:THR:OG1	2.31	0.44
2:L:33:LEU:HD13	2:L:34:ALA:N	2.33	0.44
2:L:76:ARG:O	2:L:77:ARG:NH1	2.51	0.44
1:H:50:ARG:HD2	1:H:52:TYR:CZ	2.53	0.43
2:L:188:LYS:HD3	2:L:188:LYS:HA	1.79	0.43
2:L:81:GLU:H	2:L:81:GLU:CD	2.25	0.43
2:L:83:SER:HA	2:L:104:VAL:O	2.17	0.43
3:D:143:TRP:H	3:D:143:TRP:HE3	1.65	0.42
3:D:144:LEU:O	3:D:148:VAL:HG22	2.19	0.42
3:D:177:LYS:NZ	4:D:201:EDO:H11	2.34	0.42
2:L:33:LEU:HB3	2:L:51:ALA:HB2	2.01	0.42
1:H:116:THR:HG23	1:H:147:PRO:HD2	2.02	0.42
3:D:148:VAL:HG12	3:D:179:PHE:CD1	2.55	0.41
2:L:136:LEU:HD22	2:L:175:LEU:HD22	2.03	0.41
2:L:115:VAL:HG12	2:L:207:LYS:HG3	2.02	0.41
3:D:116:TYR:HB3	3:D:175:VAL:HG12	2.02	0.40
1:H:75:LYS:HE3	1:H:75:LYS:HB3	1.75	0.40
2:L:103:LYS:NZ	5:L:402:HOH:O	2.53	0.40
2:L:195:GLU:HG3	2:L:206:THR:OG1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	226/228 (99%)	221 (98%)	5 (2%)	0	100	100
2	L	202/216 (94%)	198 (98%)	4 (2%)	0	100	100
3	C	71/100 (71%)	69 (97%)	2 (3%)	0	100	100
3	D	69/100 (69%)	69 (100%)	0	0	100	100
All	All	568/644 (88%)	557 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	191/191 (100%)	189 (99%)	2 (1%)	68	81
2	L	179/185 (97%)	178 (99%)	1 (1%)	78	88
3	C	67/91 (74%)	67 (100%)	0	100	100
3	D	64/91 (70%)	64 (100%)	0	100	100
All	All	501/558 (90%)	498 (99%)	3 (1%)	78	88

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	150	VAL
1	H	170	LEU
2	L	97	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	77	GLN
1	H	164	HIS
1	H	192	GLN
2	L	31	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	L	137	ASN
3	C	125	ASN
3	C	135	ASN
3	C	157	ASN
3	D	125	ASN
3	D	135	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	EDO	H	303	-	3,3,3	0.25	0	2,2,2	0.29	0
4	EDO	L	301	-	3,3,3	0.25	0	2,2,2	0.32	0
4	EDO	D	201	-	3,3,3	0.25	0	2,2,2	0.35	0
4	EDO	H	304	-	3,3,3	0.24	0	2,2,2	0.43	0
4	EDO	L	303	-	3,3,3	0.25	0	2,2,2	0.35	0
4	EDO	H	301	-	3,3,3	0.26	0	2,2,2	0.35	0
4	EDO	L	304	-	3,3,3	0.25	0	2,2,2	0.36	0
4	EDO	L	302	-	3,3,3	0.25	0	2,2,2	0.32	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	EDO	H	302	-	3,3,3	0.25	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	H	303	-	-	0/1/1/1	-
4	EDO	L	301	-	-	0/1/1/1	-
4	EDO	D	201	-	-	0/1/1/1	-
4	EDO	H	304	-	-	0/1/1/1	-
4	EDO	L	303	-	-	1/1/1/1	-
4	EDO	H	301	-	-	0/1/1/1	-
4	EDO	L	304	-	-	0/1/1/1	-
4	EDO	L	302	-	-	1/1/1/1	-
4	EDO	H	302	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	L	302	EDO	O1-C1-C2-O2
4	L	303	EDO	O1-C1-C2-O2
4	H	302	EDO	O1-C1-C2-O2

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	201	EDO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand EDO H 303




Ligand EDO L 301




Ligand EDO D 201




Ligand EDO H 304



Bond lengths



Bond angles




Torsions




Rings


Ligand EDO L 303




Bond lengths



Bond angles




Torsions




Rings


Ligand EDO H 301




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	H	228/228 (100%)	-0.20	3 (1%)	75	71	20, 34, 57, 80	0
2	L	208/216 (96%)	0.03	8 (3%)	44	36	20, 40, 71, 93	0
3	C	75/100 (75%)	0.48	5 (6%)	24	18	35, 55, 88, 92	0
3	D	73/100 (73%)	0.77	9 (12%)	8	7	37, 57, 96, 108	0
All	All	584/644 (90%)	0.09	25 (4%)	40	32	20, 41, 80, 108	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	143	TRP	4.4
1	H	216	CYS	3.7
3	D	138	ASP	3.4
2	L	26	SER	3.3
2	L	8	PRO	3.2
2	L	27	VAL	3.1
1	H	42	GLY	2.9
3	C	171	VAL	2.9
3	D	104	TYR	2.8
2	L	25	ALA	2.8
3	D	183	SER	2.7
3	C	162	THR	2.6
3	D	140	LEU	2.6
3	D	101	ASN	2.6
2	L	67	SER	2.5
2	L	5	THR	2.4
3	D	108	SER	2.4
2	L	93	SER	2.3
3	C	109	CYS	2.3
3	D	100	CYS	2.3
3	D	173	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	1	GLN	2.3
2	L	7	SER	2.2
3	C	173	GLN	2.1
3	C	104	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	EDO	H	303	4/4	0.67	0.18	49,55,56,74	0
4	EDO	H	301	4/4	0.76	0.20	50,53,55,58	0
4	EDO	D	201	4/4	0.82	0.21	46,49,57,61	0
4	EDO	L	302	4/4	0.84	0.14	34,41,47,59	0
4	EDO	L	304	4/4	0.89	0.12	36,48,54,55	0
4	EDO	L	301	4/4	0.90	0.13	45,48,50,53	0
4	EDO	L	303	4/4	0.91	0.13	37,45,45,56	0
4	EDO	H	304	4/4	0.93	0.12	37,39,41,44	0
4	EDO	H	302	4/4	0.93	0.12	30,32,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around EDO H 303:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around EDO H 301:


$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around EDO D 201:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)









Electron density around EDO L 302:


$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around EDO L 304:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around EDO L 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)









Electron density around EDO L 303:


$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around EDO H 304:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)








Electron density around EDO H 302:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.