



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 19, 2026 – 10:09 AM EDT

PDB ID : 9ZBX / pdb_00009zbx
Title : Crystal structure of SARS-CoV-2 receptor binding domain in complex with antibodies HB148-M4 and LY-CoV1404
Authors : Yuan, M.; Feng, Z.; Wilson, I.A.
Deposited on : 2025-11-21
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	205	9% 85% 9% 5%
2	A	222	4% 84% 11% 5%
3	B	215	6% 92% 7% .
4	H	221	% 89% 8% .
5	L	217	% 92% 6% .

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 8159 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	194	Total	C	N	O	S	0	1	0
			1544	990	259	287	8			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	531	GLY	-	expression tag	UNP P0DTC2
C	532	HIS	-	expression tag	UNP P0DTC2
C	533	HIS	-	expression tag	UNP P0DTC2
C	534	HIS	-	expression tag	UNP P0DTC2
C	535	HIS	-	expression tag	UNP P0DTC2
C	536	HIS	-	expression tag	UNP P0DTC2
C	537	HIS	-	expression tag	UNP P0DTC2

- Molecule 2 is a protein called Antibody LY-CoV1404 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	212	Total	C	N	O	S	0	0	0
			1588	1017	257	309	5			

- Molecule 3 is a protein called Antibody LY-CoV1404 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	212	Total	C	N	O	S	0	0	0
			1566	975	259	327	5			


- Molecule 4 is a protein called Antibody HB148-M4 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	215	Total	C	N	O	S	0	0	0
			1612	1021	268	317	6			

- Molecule 5 is a protein called Antibody HB148-M4 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	L	215	Total	C	N	O	S	0	1	0
			1654	1034	281	335	4			

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	O	0	0
			4	2	2		
6	C	1	Total	C	O	0	0
			4	2	2		
6	L	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	O	P	0	0
			5	4	1		

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 9 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	H	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	C	40	Total	O	0	0
			40	40		
10	A	27	Total	O	0	0
			27	27		
10	B	10	Total	O	0	0
			10	10		
10	H	45	Total	O	0	0
			45	45		
10	L	34	Total	O	0	0
			34	34		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S1




- Molecule 2: Antibody LY-CoV1404 Fab Heavy Chain




- Molecule 3: Antibody LY-CoV1404 Fab Light Chain




- Molecule 4: Antibody HB148-M4 Fab Heavy Chain



- Molecule 5: Antibody HB148-M4 Fab Light Chain





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	195.84Å 88.73Å 99.88Å 90.00° 112.00° 90.00°	Depositor
Resolution (Å)	36.14 – 2.60 36.14 – 2.60	Depositor EDS
% Data completeness (in resolution range)	86.1 (36.14-2.60) 86.4 (36.14-2.60)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.193 , 0.243 0.193 , 0.243	Depositor DCC
R_{free} test set	2109 reflections (4.28%)	wwPDB-VP
Wilson B-factor (Å ²)	43.7	Xtriage
Anisotropy	0.155	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 35.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8159	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, NAG, TRS, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.09	0/1591	0.26	0/2165
2	A	0.11	0/1629	0.32	0/2231
3	B	0.12	0/1603	0.32	0/2187
4	H	0.10	0/1649	0.30	0/2246
5	L	0.10	0/1693	0.31	0/2297
All	All	0.11	0/8165	0.30	0/11126

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1544	0	1465	11	0
2	A	1588	0	1565	16	0
3	B	1566	0	1506	9	0
4	H	1612	0	1588	11	0
5	L	1654	0	1609	9	0
6	C	8	0	12	0	0
6	L	4	0	6	0	0
7	C	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	C	14	0	13	0	0
9	H	8	0	12	0	0
10	A	27	0	0	0	0
10	B	10	0	0	0	0
10	C	40	0	0	0	0
10	H	45	0	0	0	0
10	L	34	0	0	0	0
All	All	8159	0	7776	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:188:SER:HB2	2:A:192:LYS:HG3	1.66	0.77
2:A:75:LYS:HB3	2:A:77:GLN:HG3	1.65	0.77
1:C:346:ARG:HH12	2:A:53:TRP:HE1	1.43	0.66
2:A:181:VAL:HG21	3:B:135:LEU:HD13	1.80	0.63
4:H:29:VAL:HG13	4:H:34:MET:HG3	1.81	0.63
5:L:90:GLN:NE2	5:L:93:SER:H	1.98	0.62
2:A:87:THR:HG23	2:A:110:THR:HA	1.82	0.60
2:A:67:LEU:HD12	2:A:80:LEU:HD21	1.83	0.60
3:B:105:THR:HG21	3:B:141:PRO:HB3	1.84	0.59
1:C:376:THR:HB	1:C:435:ALA:HB3	1.83	0.59
5:L:181:LEU:HD12	5:L:185:ASP:HB3	1.85	0.59
1:C:383:SER:HB3	1:C:386:LYS:HG3	1.86	0.56
4:H:85:GLU:H	4:H:85:GLU:CD	2.14	0.56
4:H:82:MET:HB3	4:H:82(C):LEU:HD21	1.89	0.55
1:C:393:THR:HA	1:C:522:ALA:HA	1.91	0.52
4:H:119:PRO:HB3	4:H:145:TYR:HB3	1.91	0.52
2:A:193:THR:HG23	2:A:210:LYS:HE3	1.91	0.51
2:A:22:CYS:HB2	2:A:36:TRP:CZ2	2.45	0.51
1:C:360:ASN:H	1:C:523:THR:HG23	1.77	0.50
4:H:20:LEU:HD12	4:H:80:LEU:HD23	1.94	0.48
4:H:23:ALA:HA	4:H:77:THR:HG22	1.95	0.48
5:L:37:GLN:HB2	5:L:47:LEU:HD11	1.95	0.48
5:L:83:PHE:CD2	5:L:106:ILE:HG13	2.49	0.47
2:A:169:VAL:HB	3:B:162:THR:HG22	1.96	0.47
4:H:83:ARG:HB2	4:H:85:GLU:OE1	2.15	0.47
3:B:83:GLU:HG3	3:B:104:LEU:O	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:36:TRP:CG	2:A:80:LEU:HD12	2.50	0.47
4:H:64:LYS:HE3	4:H:64:LYS:HB2	1.64	0.46
1:C:421:TYR:CD1	1:C:457:ARG:HB3	2.50	0.46
4:H:200:HIS:CD2	4:H:202:PRO:HD2	2.51	0.45
5:L:90:GLN:HE21	5:L:92:GLY:H	1.64	0.45
2:A:40:PRO:HB2	2:A:43:LYS:HB2	1.99	0.44
1:C:502:GLY:O	1:C:506:GLN:HG3	2.18	0.44
1:C:500:THR:HB	3:B:28:GLY:HA3	2.00	0.43
2:A:159:LEU:HD21	2:A:182:VAL:HG21	1.99	0.43
1:C:393:THR:HG21	1:C:520:ALA:HB3	2.01	0.43
5:L:83:PHE:HA	5:L:104:LEU:HD12	2.02	0.42
1:C:350:VAL:HG22	1:C:422:ASN:HB3	2.00	0.42
2:A:32:SER:HA	2:A:53:TRP:CE2	2.55	0.42
2:A:50:LEU:HD21	2:A:95:HIS:CD2	2.55	0.42
3:B:185:TRP:CE2	3:B:208:PRO:HB3	2.54	0.42
4:H:25:SER:C	4:H:27:PHE:H	2.28	0.42
1:C:378:LYS:HD2	1:C:380:TYR:CZ	2.55	0.41
2:A:13:LYS:HE2	2:A:13:LYS:HB2	1.96	0.41
4:H:100:GLY:HA2	5:L:49:TYR:CD2	2.55	0.41
3:B:188:HIS:HB2	3:B:191:TYR:CZ	2.56	0.41
2:A:66:ARG:HD2	2:A:82(A):THR:O	2.21	0.41
3:B:113:PRO:HB3	3:B:139:PHE:HB3	2.02	0.40
5:L:33:LEU:HD22	5:L:71:PHE:CD1	2.57	0.40
3:B:120:PRO:HD2	3:B:185:TRP:CZ2	2.57	0.40
5:L:113:PRO:HB3	5:L:139:PHE:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	193/205 (94%)	189 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	208/222 (94%)	204 (98%)	4 (2%)	0	100	100
3	B	210/215 (98%)	204 (97%)	6 (3%)	0	100	100
4	H	211/221 (96%)	204 (97%)	7 (3%)	0	100	100
5	L	214/217 (99%)	210 (98%)	4 (2%)	0	100	100
All	All	1036/1080 (96%)	1011 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	168/177 (95%)	168 (100%)	0	100	100
2	A	181/196 (92%)	180 (99%)	1 (1%)	78	91
3	B	177/181 (98%)	177 (100%)	0	100	100
4	H	182/186 (98%)	182 (100%)	0	100	100
5	L	186/187 (100%)	185 (100%)	1 (0%)	81	92
All	All	894/927 (96%)	892 (100%)	2 (0%)	87	95

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	192	LYS
5	L	33	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	360	ASN
1	C	414	GLN
1	C	481	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	A	16	GLN
2	A	76	ASN
2	A	171	GLN
3	B	31	ASN
4	H	81	GLN
4	H	82(A)	ASN
4	H	164	HIS
4	H	204	ASN
5	L	90	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	PO4	C	602	-	4,4,4	1.51	1 (25%)	6,6,6	0.63	0
6	EDO	C	601	-	3,3,3	0.25	0	2,2,2	0.30	0
6	EDO	L	301	-	3,3,3	0.25	0	2,2,2	0.27	0
8	NAG	C	603	1	14,14,15	0.68	0	17,19,21	1.06	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	C	604	-	3,3,3	0.25	0	2,2,2	0.29	0
9	TRS	H	301	-	7,7,7	0.33	0	9,9,9	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	C	601	-	-	0/1/1/1	-
6	EDO	L	301	-	-	1/1/1/1	-
8	NAG	C	603	1	-	3/6/23/26	0/1/1/1
6	EDO	C	604	-	-	0/1/1/1	-
9	TRS	H	301	-	-	3/9/9/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	C	602	PO4	P-O1	2.58	1.56	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	C	603	NAG	C1-O5-C5	2.70	115.80	112.19

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	C	603	NAG	C8-C7-N2-C2
8	C	603	NAG	O7-C7-N2-C2
8	C	603	NAG	O5-C5-C6-O6
9	H	301	TRS	N-C-C1-O1
9	H	301	TRS	C2-C-C1-O1
9	H	301	TRS	C3-C-C1-O1
6	L	301	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	C	194/205 (94%)	0.15	18 (9%)	14	11	18, 41, 94, 109	1 (0%)
2	A	212/222 (95%)	0.32	8 (3%)	44	38	25, 57, 87, 104	0
3	B	212/215 (98%)	0.58	13 (6%)	27	22	29, 61, 88, 109	0
4	H	215/221 (97%)	-0.19	2 (0%)	81	78	17, 41, 71, 95	0
5	L	215/217 (99%)	-0.29	3 (1%)	73	69	18, 37, 65, 75	1 (0%)
All	All	1048/1080 (97%)	0.11	44 (4%)	40	35	17, 48, 86, 109	2 (0%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	B	80	ALA	4.0
2	A	73	THR	3.6
2	A	11	LEU	3.1
3	B	209	THR	3.1
1	C	385	THR	3.0
2	A	15	THR	2.9
1	C	519	HIS	2.9
2	A	2	ILE	2.8
1	C	430	THR	2.8
2	A	9	PRO	2.7
3	B	191	TYR	2.7
1	C	381	GLY	2.7
1	C	391	CYS	2.6
1	C	368	LEU	2.6
3	B	146	VAL	2.6
2	A	10	THR	2.6
3	B	156	LYS	2.5
1	C	527	PRO	2.5
1	C	525	CYS	2.5
1	C	372	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	522	ALA	2.5
3	B	186	LYS	2.4
3	B	190	SER	2.4
5	L	95	PRO	2.3
1	C	365	TYR	2.3
3	B	155	VAL	2.3
1	C	373	SER	2.3
4	H	190	GLY	2.2
3	B	207	ALA	2.2
3	B	13	GLY	2.2
1	C	382	VAL	2.2
3	B	144	VAL	2.2
1	C	371	SER	2.2
5	L	94	SER	2.2
2	A	102	HIS	2.1
3	B	56	SER	2.1
4	H	129	LYS	2.1
1	C	518	LEU	2.1
1	C	334	ASN	2.1
2	A	23	THR	2.0
1	C	335	LEU	2.0
5	L	213	GLU	2.0
3	B	179	SER	2.0
1	C	517	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	EDO	L	301	4/4	0.68	0.25	55,59,63,67	0
8	NAG	C	603	14/15	0.77	0.14	67,79,93,95	0
9	TRS	H	301	8/8	0.85	0.23	46,56,70,80	0
6	EDO	C	604	4/4	0.88	0.14	39,44,47,53	0
6	EDO	C	601	4/4	0.97	0.11	42,43,45,50	0
7	PO4	C	602	5/5	0.98	0.07	35,36,42,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.