



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 19, 2026 – 10:10 AM EDT

PDB ID : 9ZBY / pdb_00009zby
Title : Crystal structure of SARS-CoV-2 Omicron BA.1 receptor binding domain in complex with antibodies HB148-M4 and LY-CoV1404
Authors : Feng, Z.; Yuan, M.; Wilson, I.A.
Deposited on : 2025-11-21
Resolution : 2.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1819 (2.76-2.72)
Clashscore	190562	1866 (2.76-2.72)
Ramachandran outliers	187476	1830 (2.76-2.72)
Sidechain outliers	187428	1831 (2.76-2.72)
RSRZ outliers	180081	1819 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	221	81% 17% .
2	L	217	95% . .
3	A	222	4% 83% 13% .
4	B	215	11% 82% 16% .
5	C	205	7% 82% 10% . 7%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 8052 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antibody HB148-M4 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	215	Total	C	N	O	S	0	0	0
			1610	1020	268	316	6			

- Molecule 2 is a protein called Antibody HB148-M4 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	215	Total	C	N	O	S	0	1	0
			1654	1034	281	335	4			

- Molecule 3 is a protein called Antibody LY-CoV1404 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	213	Total	C	N	O	S	0	0	0
			1608	1027	263	313	5			

- Molecule 4 is a protein called Antibody LY-CoV1404 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	212	Total	C	N	O	S	0	0	0
			1566	975	259	327	5			

- Molecule 5 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	C	190	Total	C	N	O	S	0	0	0
			1522	980	258	276	8			

There are 25 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	339	ASP	GLY	conflict	UNP P0DTC2
C	371	LEU	SER	conflict	UNP P0DTC2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	373	PRO	SER	conflict	UNP P0DTC2
C	375	PHE	SER	conflict	UNP P0DTC2
C	417	ASN	LYS	conflict	UNP P0DTC2
C	440	LYS	ASN	conflict	UNP P0DTC2
C	446	SER	GLY	conflict	UNP P0DTC2
C	477	ASN	SER	conflict	UNP P0DTC2
C	478	LYS	THR	conflict	UNP P0DTC2
C	484	ALA	GLU	conflict	UNP P0DTC2
C	493	ARG	GLN	conflict	UNP P0DTC2
C	496	SER	GLY	conflict	UNP P0DTC2
C	498	ARG	GLN	conflict	UNP P0DTC2
C	501	TYR	ASN	conflict	UNP P0DTC2
C	505	HIS	TYR	conflict	UNP P0DTC2
C	528	LYS	-	expression tag	UNP P0DTC2
C	529	LYS	-	expression tag	UNP P0DTC2
C	530	SER	-	expression tag	UNP P0DTC2
C	531	GLY	-	expression tag	UNP P0DTC2
C	532	HIS	-	expression tag	UNP P0DTC2
C	533	HIS	-	expression tag	UNP P0DTC2
C	534	HIS	-	expression tag	UNP P0DTC2
C	535	HIS	-	expression tag	UNP P0DTC2
C	536	HIS	-	expression tag	UNP P0DTC2
C	537	HIS	-	expression tag	UNP P0DTC2

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	H	1	Total C O 4 2 2	0	0
6	L	1	Total C O 4 2 2	0	0
6	L	1	Total C O 4 2 2	0	0
6	L	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0

- Molecule 7 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	L	1	Total O P 5 4 1	0	0
7	C	1	Total O P 5 4 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	H	12	Total 12	O 12	0	0
8	L	10	Total 10	O 10	0	0
8	A	8	Total 8	O 8	0	0
8	B	6	Total 6	O 6	0	0
8	C	14	Total 14	O 14	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Antibody HB148-M4 Fab Heavy Chain




- Molecule 2: Antibody HB148-M4 Fab Light Chain




- Molecule 3: Antibody LY-CoV1404 Fab Heavy Chain






- Molecule 4: Antibody LY-CoV1404 Fab Light Chain





● Molecule 5: Spike protein S1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	196.86Å 88.03Å 100.81Å 90.00° 110.27° 90.00°	Depositor
Resolution (Å)	40.75 – 2.73 40.75 – 2.73	Depositor EDS
% Data completeness (in resolution range)	78.3 (40.75-2.73) 79.9 (40.75-2.73)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.58 (at 2.73Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.200 , 0.254 0.200 , 0.254	Depositor DCC
R_{free} test set	2000 reflections (4.52%)	wwPDB-VP
Wilson B-factor (Å ²)	43.5	Xtriage
Anisotropy	0.279	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8052	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.10	0/1647	0.27	0/2243
2	L	0.09	0/1693	0.27	0/2297
3	A	0.12	0/1649	0.29	0/2255
4	B	0.10	0/1603	0.30	0/2187
5	C	0.09	0/1567	0.26	0/2131
All	All	0.10	0/8159	0.28	0/11113

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1610	0	1586	21	0
2	L	1654	0	1609	5	0
3	A	1608	0	1612	21	0
4	B	1566	0	1506	23	0
5	C	1522	0	1449	12	0
6	C	16	0	24	0	0
6	H	4	0	6	0	0
6	L	12	0	18	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	C	5	0	0	0	0
7	L	5	0	0	0	0
8	A	8	0	0	0	0
8	B	6	0	0	0	0
8	C	14	0	0	0	0
8	H	12	0	0	0	0
8	L	10	0	0	0	0
All	All	8052	0	7810	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:91:TYR:HB3	2:L:96:TYR:HB3	1.70	0.72
5:C:411:ALA:HB3	5:C:414:GLN:HG3	1.74	0.68
3:A:29:LEU:HD13	3:A:53:TRP:HA	1.77	0.67
2:L:37:GLN:HB2	2:L:47:LEU:HD11	1.76	0.66
1:H:51:ILE:HG13	1:H:57:THR:HG22	1.79	0.64
3:A:39:GLN:HE22	4:B:38:GLN:HE22	1.45	0.63
4:B:80:ALA:HA	4:B:106:VAL:HG11	1.81	0.63
1:H:66:ARG:NH2	1:H:86:ASP:OD2	2.32	0.62
1:H:197:ASN:ND2	1:H:208:ASP:OD2	2.33	0.62
3:A:29:LEU:HD12	3:A:71:LYS:CD	2.30	0.61
1:H:29:VAL:HG13	1:H:34:MET:HG3	1.82	0.61
3:A:5:LYS:HG2	3:A:23:THR:HB	1.84	0.59
4:B:60:ASN:N	4:B:60:ASN:OD1	2.33	0.59
4:B:37:GLN:HB3	4:B:47:MET:HE3	1.84	0.59
3:A:181:VAL:HG21	4:B:135:LEU:HD13	1.86	0.57
4:B:166:LYS:HA	4:B:172:TYR:HD2	1.69	0.57
4:B:181:THR:HG23	4:B:184:GLN:H	1.70	0.56
1:H:7:SER:OG	1:H:21:SER:OG	2.25	0.54
1:H:35:SER:OG	1:H:95:ASP:OD2	2.22	0.54
1:H:87:THR:HG23	1:H:110:THR:HA	1.90	0.53
5:C:449:TYR:OH	5:C:498:ARG:NH2	2.42	0.53
3:A:29:LEU:HD12	3:A:71:LYS:HD2	1.91	0.52
3:A:32:SER:HA	3:A:53:TRP:CD2	2.45	0.52
3:A:6:GLU:N	3:A:6:GLU:OE2	2.43	0.51
3:A:12:VAL:HG23	3:A:111:VAL:HG22	1.91	0.51
3:A:119:PRO:HB3	3:A:145:TYR:HB3	1.93	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:103:TRP:HD1	6:L:302:EDO:H11	1.75	0.51
4:B:150:ALA:HB1	4:B:188:HIS:HD2	1.76	0.50
1:H:34:MET:HB3	1:H:78:LEU:HD22	1.94	0.49
4:B:181:THR:HG22	4:B:184:GLN:HB2	1.95	0.49
1:H:200:HIS:ND1	1:H:203:SER:OG	2.44	0.49
3:A:29:LEU:HD12	3:A:71:LYS:HD3	1.94	0.48
3:A:84:PRO:HA	3:A:111:VAL:HB	1.95	0.48
1:H:119:PRO:HB3	1:H:145:TYR:HB3	1.95	0.48
4:B:185:TRP:HH2	4:B:206:VAL:HG12	1.79	0.48
2:L:67:SER:HB3	4:B:93:THR:HG21	1.95	0.47
5:C:365:TYR:CE2	5:C:387:LEU:HB3	2.49	0.47
3:A:200:HIS:CD2	3:A:202:PRO:HD2	2.48	0.47
3:A:210:LYS:HE2	3:A:212:GLU:HG2	1.95	0.47
5:C:497:PHE:CE2	5:C:507:PRO:HB3	2.50	0.47
1:H:195:ILE:HG13	1:H:210:LYS:HG3	1.97	0.46
4:B:49:PHE:CE2	4:B:55:PRO:HG3	2.51	0.46
4:B:196:THR:HG1	4:B:201:THR:HG1	1.50	0.46
1:H:59:TYR:HB2	1:H:64:LYS:HD2	1.97	0.46
1:H:22:CYS:HB3	1:H:78:LEU:HB3	1.97	0.46
5:C:382:VAL:HB	5:C:387:LEU:HD11	1.98	0.46
5:C:408:ARG:NH2	5:C:414:GLN:OE1	2.49	0.45
4:B:46:LEU:HD21	4:B:49:PHE:HD2	1.81	0.45
4:B:148:TRP:CE3	4:B:178:LEU:HD12	2.52	0.45
5:C:501:TYR:O	5:C:506:GLN:NE2	2.38	0.45
1:H:13:GLN:H	1:H:13:GLN:CD	2.25	0.44
3:A:119:PRO:HD3	3:A:200:HIS:ND1	2.33	0.44
4:B:144:VAL:HG23	4:B:195:VAL:HG13	1.99	0.44
4:B:27(B):ASP:N	4:B:27(B):ASP:OD1	2.50	0.44
1:H:82:MET:HE2	1:H:82(C):LEU:HD21	1.99	0.43
5:C:388:ASN:OD1	5:C:388:ASN:N	2.51	0.43
3:A:124:LEU:HB3	4:B:118:PHE:CD1	2.53	0.43
1:H:8:GLY:O	1:H:18:LEU:HD21	2.17	0.43
1:H:12:ILE:HD13	1:H:82(C):LEU:HD12	1.99	0.43
3:A:168:ALA:HA	3:A:178:LEU:HB3	2.00	0.43
3:A:94:HIS:ND1	3:A:101:ASP:OD2	2.48	0.42
1:H:25:SER:C	1:H:27:PHE:H	2.28	0.42
4:B:166:LYS:HA	4:B:172:TYR:CD2	2.53	0.42
5:C:490:PHE:CE2	5:C:492:LEU:HB2	2.53	0.42
1:H:53:PRO:HA	1:H:71:ARG:HH12	1.83	0.42
3:A:191:THR:O	3:A:192:LYS:HD2	2.20	0.41
2:L:32:TYR:HA	2:L:91:TYR:CE1	2.54	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:26:GLU:OE1	5:C:478:LYS:NZ	2.51	0.41
5:C:377:PHE:CZ	5:C:384:PRO:HG3	2.55	0.41
4:B:81:GLU:OE1	4:B:81:GLU:N	2.54	0.41
2:L:31:SER:HA	2:L:50:GLY:HA2	2.03	0.41
4:B:130:ALA:HB3	4:B:180:LEU:HD12	2.03	0.41
4:B:81:GLU:H	4:B:81:GLU:CD	2.29	0.40
3:A:1:GLN:OE1	3:A:1:GLN:N	2.46	0.40
3:A:56:ASP:OD2	5:C:444:LYS:NZ	2.54	0.40
4:B:128:ASN:HA	4:B:182:PRO:HG2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	211/221 (96%)	205 (97%)	6 (3%)	0	100	100
2	L	214/217 (99%)	207 (97%)	7 (3%)	0	100	100
3	A	209/222 (94%)	206 (99%)	3 (1%)	0	100	100
4	B	210/215 (98%)	201 (96%)	9 (4%)	0	100	100
5	C	186/205 (91%)	182 (98%)	4 (2%)	0	100	100
All	All	1030/1080 (95%)	1001 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	181/186 (97%)	180 (99%)	1 (1%)	78	87
2	L	186/187 (100%)	186 (100%)	0	100	100
3	A	188/196 (96%)	187 (100%)	1 (0%)	81	88
4	B	177/181 (98%)	173 (98%)	4 (2%)	44	65
5	C	165/179 (92%)	160 (97%)	5 (3%)	36	58
All	All	897/929 (97%)	886 (99%)	11 (1%)	63	77

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	99	VAL
3	A	151	THR
4	B	29	ASP
4	B	60	ASN
4	B	75	ILE
4	B	155	VAL
5	C	383	SER
5	C	387	LEU
5	C	388	ASN
5	C	389	ASP
5	C	483	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	L	147	GLN
3	A	171	GLN
4	B	38	GLN
4	B	188	HIS
5	C	448	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	EDO	L	304	-	3,3,3	0.26	0	2,2,2	0.21	0
6	EDO	C	602	-	3,3,3	0.25	0	2,2,2	0.25	0
7	PO4	L	301	-	4,4,4	1.58	1 (25%)	6,6,6	0.48	0
6	EDO	C	604	-	3,3,3	0.26	0	2,2,2	0.34	0
6	EDO	C	605	-	3,3,3	0.25	0	2,2,2	0.26	0
6	EDO	C	603	-	3,3,3	0.25	0	2,2,2	0.34	0
6	EDO	L	303	-	3,3,3	0.26	0	2,2,2	0.29	0
7	PO4	C	601	-	4,4,4	1.58	1 (25%)	6,6,6	0.54	0
6	EDO	L	302	-	3,3,3	0.24	0	2,2,2	0.30	0
6	EDO	H	301	-	3,3,3	0.25	0	2,2,2	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	L	304	-	-	0/1/1/1	-
6	EDO	C	602	-	-	1/1/1/1	-
6	EDO	C	604	-	-	0/1/1/1	-
6	EDO	C	605	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	C	603	-	-	0/1/1/1	-
6	EDO	L	303	-	-	1/1/1/1	-
6	EDO	L	302	-	-	0/1/1/1	-
6	EDO	H	301	-	-	1/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	L	301	PO4	P-O1	2.76	1.57	1.50
7	C	601	PO4	P-O1	2.75	1.57	1.50

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	H	301	EDO	O1-C1-C2-O2
6	L	303	EDO	O1-C1-C2-O2
6	C	602	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	L	302	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	215/221 (97%)	-0.12	2 (0%) 81 81	13, 37, 65, 95	0
2	L	215/217 (99%)	-0.22	1 (0%) 87 87	11, 30, 61, 94	1 (0%)
3	A	213/222 (95%)	0.55	9 (4%) 40 39	29, 58, 84, 98	0
4	B	212/215 (98%)	1.00	23 (10%) 11 10	23, 74, 96, 107	0
5	C	190/205 (92%)	0.34	15 (7%) 18 19	14, 39, 100, 112	0
All	All	1045/1080 (96%)	0.31	50 (4%) 35 35	11, 48, 90, 112	1 (0%)

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	C	374	PHE	4.6
4	B	47	MET	3.8
5	C	367	VAL	3.7
3	A	10	THR	3.6
5	C	373	PRO	3.4
3	A	27	PHE	3.3
2	L	1	GLU	3.1
5	C	372	ALA	3.0
5	C	383	SER	3.0
4	B	209	THR	2.9
4	B	75	ILE	2.8
3	A	140	CYS	2.8
5	C	375	PHE	2.7
5	C	527	PRO	2.7
4	B	82	ASP	2.7
4	B	159	VAL	2.7
4	B	144	VAL	2.7
4	B	155	VAL	2.6
3	A	101	ASP	2.6
5	C	519	HIS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	B	143	ALA	2.6
3	A	15	THR	2.6
4	B	46	LEU	2.5
4	B	185	TRP	2.5
1	H	129	LYS	2.5
4	B	186	LYS	2.5
4	B	16	GLY	2.5
5	C	381	GLY	2.5
1	H	1	GLU	2.5
5	C	336	CYS	2.4
4	B	62	PHE	2.4
4	B	80	ALA	2.4
4	B	27	SER	2.4
4	B	2	SER	2.3
5	C	362	VAL	2.3
4	B	106(A)	LEU	2.3
3	A	212	GLU	2.3
4	B	76	SER	2.3
4	B	11	VAL	2.3
4	B	206	VAL	2.3
3	A	98	SER	2.3
4	B	188	HIS	2.2
3	A	120	SER	2.2
4	B	30	TYR	2.1
5	C	382	VAL	2.1
5	C	338	PHE	2.1
5	C	442	ASP	2.0
3	A	167	PRO	2.0
4	B	141	PRO	2.0
5	C	337	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EDO	L	303	4/4	0.80	0.14	49,49,56,63	0
7	PO4	L	301	5/5	0.80	0.15	60,88,133,138	0
6	EDO	C	603	4/4	0.83	0.20	36,48,53,53	0
6	EDO	L	302	4/4	0.84	0.15	40,42,48,50	0
6	EDO	C	602	4/4	0.84	0.23	40,43,44,86	0
6	EDO	C	605	4/4	0.85	0.23	45,46,52,58	0
6	EDO	L	304	4/4	0.90	0.20	35,39,41,66	0
6	EDO	C	604	4/4	0.91	0.23	37,38,49,53	0
6	EDO	H	301	4/4	0.92	0.10	25,34,38,45	0
7	PO4	C	601	5/5	0.95	0.10	29,37,54,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.