



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 20, 2026 – 12:05 PM EDT

PDB ID : 9ZM5 / pdb_00009zm5
Title : Crystal structure of the Arabidopsis NPR1-NIMIN1 complex
Authors : Zhang, S.; Gish, M.L.; Zheng, N.
Deposited on : 2025-12-09
Resolution : 3.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1434 (3.38-3.30)
Clashscore	190562	1479 (3.38-3.30)
Ramachandran outliers	187476	1456 (3.38-3.30)
Sidechain outliers	187428	1455 (3.38-3.30)
RSRZ outliers	180081	1434 (3.38-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	529	
1	C	529	
2	B	42	
2	D	42	
3	M	4	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	N	4	 100%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8348 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltodextrin-binding protein,Regulatory protein NPR1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	500	Total	C	N	O	S	0	0	0
			3856	2463	638	740	15			
1	C	500	Total	C	N	O	S	0	0	0
			3856	2463	638	740	15			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	108	ALA	ASP	conflict	UNP A0AAX4LAZ4
A	109	ALA	LYS	conflict	UNP A0AAX4LAZ4
A	198	ALA	GLU	conflict	UNP A0AAX4LAZ4
A	199	ALA	ASN	conflict	UNP A0AAX4LAZ4
A	265	ALA	LYS	conflict	UNP A0AAX4LAZ4
A	388	ALA	LYS	conflict	UNP A0AAX4LAZ4
A	389	ALA	ASP	conflict	UNP A0AAX4LAZ4
A	393	ASN	-	linker	UNP A0AAX4LAZ4
A	394	ALA	-	linker	UNP A0AAX4LAZ4
A	395	ALA	-	linker	UNP A0AAX4LAZ4
A	396	ALA	-	linker	UNP A0AAX4LAZ4
A	397	ALA	-	linker	UNP A0AAX4LAZ4
A	398	ALA	-	linker	UNP A0AAX4LAZ4
A	444	GLY	-	linker	UNP P93002
A	445	SER	-	linker	UNP P93002
A	446	GLY	-	linker	UNP P93002
A	447	SER	-	linker	UNP P93002
A	448	GLY	-	linker	UNP P93002
C	108	ALA	ASP	conflict	UNP A0AAX4LAZ4
C	109	ALA	LYS	conflict	UNP A0AAX4LAZ4
C	198	ALA	GLU	conflict	UNP A0AAX4LAZ4
C	199	ALA	ASN	conflict	UNP A0AAX4LAZ4
C	265	ALA	LYS	conflict	UNP A0AAX4LAZ4
C	388	ALA	LYS	conflict	UNP A0AAX4LAZ4
C	389	ALA	ASP	conflict	UNP A0AAX4LAZ4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	393	ASN	-	linker	UNP A0AAX4LAZ4
C	394	ALA	-	linker	UNP A0AAX4LAZ4
C	395	ALA	-	linker	UNP A0AAX4LAZ4
C	396	ALA	-	linker	UNP A0AAX4LAZ4
C	397	ALA	-	linker	UNP A0AAX4LAZ4
C	398	ALA	-	linker	UNP A0AAX4LAZ4
C	446	GLY	-	linker	UNP P93002
C	447	SER	-	linker	UNP P93002
C	448	GLY	-	linker	UNP P93002
C	449	SER	-	linker	UNP P93002
C	450	GLY	-	linker	UNP P93002

- Molecule 2 is a protein called Protein NIM1-INTERACTING 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	30	Total	C	N	O	0	0	0
			270	169	51	50			
2	D	31	Total	C	N	O	0	0	0
			276	172	52	52			

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	M	4	Total	C	O	0	0	0
			45	24	21			
3	N	4	Total	C	O	0	0	0
			45	24	21			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltodextrin-binding protein,Regulatory protein NPR1



- Molecule 1: Maltodextrin-binding protein,Regulatory protein NPR1



- Molecule 2: Protein NIM1-INTERACTING 1



- Molecule 2: Protein NIM1-INTERACTING 1



- Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



GLC1
GLC2
GLC3
GLC4

- Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain N:

100%

GLC1
GLC2
GLC3
GLC4

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	118.48Å 144.93Å 145.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	3.50 – 3.33 3.50 – 3.33	Depositor EDS
% Data completeness (in resolution range)	96.6 (3.50-3.33) 12.4 (3.50-3.33)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 3.32Å)	Xtrriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.228 , 0.261 0.228 , 0.263	Depositor DCC
R_{free} test set	823 reflections (4.41%)	wwPDB-VP
Wilson B-factor (Å ²)	81.0	Xtrriage
Anisotropy	0.254	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.03 , 0.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.71	EDS
Total number of atoms	8348	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 34.90 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.3429e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.17	1/3935 (0.0%)	1.21	35/5328 (0.7%)
1	C	1.23	2/3935 (0.1%)	1.21	38/5328 (0.7%)
2	B	1.13	0/273	1.17	3/359 (0.8%)
2	D	1.18	0/279	1.15	1/367 (0.3%)
All	All	1.20	3/8422 (0.0%)	1.21	77/11382 (0.7%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	90	HIS	CB-CG	-5.96	1.41	1.50
1	C	90	HIS	CE1-NE2	-5.45	1.27	1.32
1	A	288	LEU	CA-C	-5.01	1.46	1.52

All (77) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	448	GLY	N-CA-C	-13.32	96.22	112.33
1	C	424	PHE	CA-CB-CG	8.68	122.48	113.80
1	A	424	PHE	CA-C-N	7.53	127.54	119.78
1	A	424	PHE	C-N-CA	7.53	127.54	119.78
1	A	82	GLY	CA-C-N	7.53	127.69	120.31
1	A	82	GLY	C-N-CA	7.53	127.69	120.31
1	A	96	TYR	N-CA-C	-7.47	103.78	113.12
1	C	254	GLY	CA-C-N	7.43	128.09	119.47
1	C	254	GLY	C-N-CA	7.43	128.09	119.47
1	A	254	GLY	CA-C-N	7.35	128.00	119.47
1	A	254	GLY	C-N-CA	7.35	128.00	119.47
1	C	179	GLU	CA-C-N	7.23	126.93	119.56
1	C	179	GLU	C-N-CA	7.23	126.93	119.56
1	C	323	LYS	CA-C-N	7.19	127.23	119.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	323	LYS	C-N-CA	7.19	127.23	119.90
1	C	424	PHE	CA-C-N	7.04	127.45	119.92
1	C	424	PHE	C-N-CA	7.04	127.45	119.92
1	C	448	GLY	CA-C-O	-7.00	116.99	122.52
1	C	82	GLY	CA-C-N	6.97	127.14	120.31
1	C	82	GLY	C-N-CA	6.97	127.14	120.31
1	C	151	PRO	CA-C-N	6.78	126.74	120.03
1	C	151	PRO	C-N-CA	6.78	126.74	120.03
1	A	151	PRO	CA-C-N	6.77	126.73	120.03
1	A	151	PRO	C-N-CA	6.77	126.73	120.03
1	A	476	CYS	N-CA-C	-6.48	105.02	113.12
1	C	106	THR	CA-C-N	6.42	126.34	119.85
1	C	106	THR	C-N-CA	6.42	126.34	119.85
1	C	96	TYR	N-CA-C	-6.37	105.53	113.55
1	A	359	ILE	CA-C-N	6.08	127.44	119.84
1	A	359	ILE	C-N-CA	6.08	127.44	119.84
1	C	526	ASP	CA-CB-CG	6.04	118.64	112.60
1	A	273	LEU	CA-C-N	6.03	125.99	119.78
1	A	273	LEU	C-N-CA	6.03	125.99	119.78
2	D	45	LYS	N-CA-C	-5.96	105.09	112.90
1	C	343	ILE	N-CA-C	-5.96	104.51	111.00
2	B	43	GLU	N-CA-C	5.85	118.82	110.10
2	B	52	LEU	N-CA-C	-5.82	104.86	111.14
1	A	179	GLU	CA-C-N	5.80	125.48	119.56
1	A	179	GLU	C-N-CA	5.80	125.48	119.56
1	A	106	THR	CA-C-N	5.79	125.70	119.85
1	A	106	THR	C-N-CA	5.79	125.70	119.85
1	A	58	GLY	N-CA-C	-5.76	101.85	112.62
1	C	359	ILE	CA-C-N	5.69	126.07	119.47
1	C	359	ILE	C-N-CA	5.69	126.07	119.47
1	C	356	MET	CA-C-N	5.69	125.64	119.78
1	C	356	MET	C-N-CA	5.69	125.64	119.78
1	A	229	HIS	N-CA-C	-5.65	106.52	113.41
1	A	271	THR	N-CA-C	5.64	115.73	108.34
1	C	273	LEU	CA-C-N	5.56	125.50	119.78
1	C	273	LEU	C-N-CA	5.56	125.50	119.78
1	C	119	THR	N-CA-C	-5.54	106.36	113.23
1	A	473	PHE	CA-C-N	5.48	125.14	119.56
1	A	473	PHE	C-N-CA	5.48	125.14	119.56
1	C	475	PHE	CA-C-N	5.37	124.98	119.56
1	C	475	PHE	C-N-CA	5.37	124.98	119.56
1	A	356	MET	CA-C-N	5.31	125.25	119.78

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	356	MET	C-N-CA	5.31	125.25	119.78
1	A	44	ASN	N-CA-C	-5.27	105.21	111.69
2	B	46	ILE	N-CA-C	5.25	116.15	111.90
1	C	352	LYS	N-CA-C	-5.23	107.03	113.41
1	A	343	ILE	N-CA-C	-5.23	105.30	111.00
1	A	119	THR	N-CA-C	-5.22	106.76	113.23
1	C	208	GLY	CA-C-N	-5.17	115.65	122.16
1	C	208	GLY	C-N-CA	-5.17	115.65	122.16
1	A	49	VAL	N-CA-C	-5.14	105.53	110.72
1	A	447	SER	CA-C-N	5.13	130.94	121.70
1	A	447	SER	C-N-CA	5.13	130.94	121.70
1	A	87	PHE	CA-CB-CG	5.13	118.93	113.80
1	C	424	PHE	CB-CA-C	5.09	116.69	108.91
1	A	327	ALA	N-CA-C	-5.07	103.70	110.55
1	C	148	LEU	CA-C-N	5.07	125.34	119.47
1	C	148	LEU	C-N-CA	5.07	125.34	119.47
1	C	84	ASP	N-CA-C	-5.05	105.48	111.69
1	A	208	GLY	CA-C-N	-5.04	115.81	122.16
1	A	208	GLY	C-N-CA	-5.04	115.81	122.16
1	C	296	SER	CA-C-N	5.03	125.47	120.04
1	C	296	SER	C-N-CA	5.03	125.47	120.04

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3856	0	3829	19	2
1	C	3856	0	3829	18	0
2	B	270	0	271	0	0
2	D	276	0	276	2	1
3	M	45	0	39	0	0
3	N	45	0	39	0	0
All	All	8348	0	8283	37	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:407:LYS:HE2	1:A:446:GLY:CA	1.35	1.55
1:A:407:LYS:CE	1:A:446:GLY:HA2	1.67	1.24
1:C:407:LYS:HZ3	1:C:448:GLY:N	1.45	1.13
1:A:407:LYS:CE	1:A:446:GLY:CA	2.30	1.02
1:A:407:LYS:HE2	1:A:446:GLY:HA3	1.39	1.01
1:C:407:LYS:NZ	1:C:448:GLY:H	1.63	0.96
1:C:407:LYS:NZ	1:C:448:GLY:CA	2.29	0.95
1:C:407:LYS:NZ	1:C:448:GLY:N	2.15	0.94
1:C:407:LYS:NZ	1:C:448:GLY:HA2	1.83	0.94
1:C:407:LYS:HZ3	1:C:448:GLY:CA	1.84	0.90
1:C:407:LYS:HZ3	1:C:448:GLY:H	1.16	0.87
1:A:407:LYS:HE3	1:A:455:HIS:CE1	2.10	0.86
1:A:407:LYS:HE2	1:A:446:GLY:HA2	0.78	0.77
1:C:407:LYS:HE2	1:C:457:HIS:CE1	2.20	0.76
1:C:407:LYS:HZ2	1:C:448:GLY:CA	2.02	0.69
1:C:407:LYS:HZ2	1:C:448:GLY:HA2	1.55	0.68
1:A:407:LYS:CE	1:A:455:HIS:CE1	2.78	0.66
1:A:407:LYS:CE	1:A:446:GLY:HA3	2.13	0.66
2:D:47:ASP:OD2	2:D:51:LYS:NZ	2.37	0.56
1:C:407:LYS:HD2	2:D:63:ARG:NH2	2.21	0.55
1:A:52:LYS:NZ	1:A:314:GLU:OE1	2.40	0.54
1:A:407:LYS:NZ	1:A:450:ARG:O	2.42	0.51
1:A:114:LYS:NZ	1:C:469:GLU:OE1	2.44	0.50
1:C:57:THR:OG1	1:C:58:GLY:N	2.44	0.50
1:A:499:THR:OG1	1:A:500:ALA:N	2.45	0.50
1:C:235:ASP:OD1	1:C:235:ASP:C	2.55	0.48
1:C:407:LYS:CE	1:C:448:GLY:HA2	2.43	0.47
1:A:210:ASP:C	1:A:210:ASP:OD1	2.56	0.45
1:C:407:LYS:CE	1:C:457:HIS:CE1	2.95	0.45
1:A:407:LYS:NZ	1:A:455:HIS:CE1	2.85	0.45
1:A:235:ASP:OD1	1:A:235:ASP:C	2.57	0.44
1:A:271:THR:OG1	1:A:272:VAL:N	2.51	0.43
1:A:359:ILE:HA	1:A:360:PRO:HD3	1.93	0.42
1:C:501:THR:OG1	1:C:502:ALA:N	2.52	0.42
1:C:184:TRP:N	1:C:185:PRO:CD	2.83	0.41
1:A:415:ASN:C	1:A:415:ASN:HD22	2.28	0.41
1:A:424:PHE:HA	1:A:425:PRO:HD3	1.93	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:233:ASP:OD1	1:A:233:ASP:OD1[3_555]	1.65	0.55
1:A:447:SER:OG	2:D:62:ARG:NE[6_445]	1.91	0.29

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	496/529 (94%)	489 (99%)	4 (1%)	3 (1%)	21	51
1	C	496/529 (94%)	484 (98%)	11 (2%)	1 (0%)	43	71
2	B	28/42 (67%)	27 (96%)	0	1 (4%)	2	18
2	D	29/42 (69%)	28 (97%)	1 (3%)	0	100	100
All	All	1049/1142 (92%)	1028 (98%)	16 (2%)	5 (0%)	24	55

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	447	SER
2	B	44	LYS
1	A	28	ILE
1	C	29	GLU
1	A	59	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	394/415 (95%)	394 (100%)	0	100	100
1	C	394/415 (95%)	394 (100%)	0	100	100
2	B	28/39 (72%)	28 (100%)	0	100	100
2	D	29/39 (74%)	29 (100%)	0	100	100
All	All	845/908 (93%)	845 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	455	HIS
1	C	65	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	GLC	M	1	3	12,12,12	0.90	0	17,17,17	1.24	1 (5%)
3	GLC	M	2	3	11,11,12	1.40	2 (18%)	15,15,17	1.60	1 (6%)
3	GLC	M	3	3	11,11,12	1.37	3 (27%)	15,15,17	2.32	4 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GLC	M	4	3	11,11,12	1.57	2 (18%)	15,15,17	2.61	2 (13%)
3	GLC	N	1	3	12,12,12	0.92	0	17,17,17	1.15	1 (5%)
3	GLC	N	2	3	11,11,12	1.11	1 (9%)	15,15,17	1.71	1 (6%)
3	GLC	N	3	3	11,11,12	1.27	1 (9%)	15,15,17	1.58	2 (13%)
3	GLC	N	4	3	11,11,12	1.33	3 (27%)	15,15,17	2.78	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	M	1	3	-	2/2/22/22	0/1/1/1
3	GLC	M	2	3	-	0/2/19/22	0/1/1/1
3	GLC	M	3	3	-	2/2/19/22	0/1/1/1
3	GLC	M	4	3	-	1/2/19/22	0/1/1/1
3	GLC	N	1	3	-	2/2/22/22	0/1/1/1
3	GLC	N	2	3	-	0/2/19/22	0/1/1/1
3	GLC	N	3	3	-	2/2/19/22	0/1/1/1
3	GLC	N	4	3	-	2/2/19/22	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	M	4	GLC	C2-C3	3.28	1.57	1.52
3	M	4	GLC	C1-C2	2.75	1.58	1.52
3	N	3	GLC	C2-C3	2.68	1.56	1.52
3	M	3	GLC	C2-C3	2.62	1.56	1.52
3	M	2	GLC	C2-C3	2.61	1.56	1.52
3	N	4	GLC	O5-C1	2.33	1.47	1.43
3	N	2	GLC	C4-C5	2.28	1.57	1.53
3	M	2	GLC	C4-C5	2.28	1.57	1.53
3	M	3	GLC	O4-C4	2.12	1.48	1.43
3	N	4	GLC	O5-C5	2.09	1.47	1.43
3	N	4	GLC	C2-C3	2.05	1.55	1.52
3	M	3	GLC	C4-C5	2.04	1.57	1.53

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	N	4	GLC	C1-O5-C5	10.02	125.61	112.19
3	M	4	GLC	C1-O5-C5	8.99	124.23	112.19
3	N	2	GLC	C1-O5-C5	6.32	120.66	112.19
3	M	3	GLC	C1-O5-C5	5.85	120.03	112.19
3	M	2	GLC	C1-O5-C5	5.78	119.93	112.19
3	N	3	GLC	C1-O5-C5	5.19	119.15	112.19
3	M	3	GLC	O4-C4-C5	4.05	119.30	109.32
3	M	4	GLC	C1-C2-C3	3.48	114.71	109.64
3	N	1	GLC	C1-O5-C5	3.05	119.54	113.65
3	M	1	GLC	C1-O5-C5	2.99	119.44	113.65
3	M	3	GLC	O5-C5-C6	-2.73	102.34	107.66
3	M	3	GLC	O2-C2-C1	-2.34	103.88	109.22
3	N	3	GLC	O2-C2-C1	-2.13	104.35	109.22

There are no chirality outliers.


All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	M	3	GLC	O5-C5-C6-O6
3	M	1	GLC	O5-C5-C6-O6
3	M	3	GLC	C4-C5-C6-O6
3	N	1	GLC	O5-C5-C6-O6
3	M	1	GLC	C4-C5-C6-O6
3	N	4	GLC	C4-C5-C6-O6
3	M	4	GLC	O5-C5-C6-O6
3	N	4	GLC	O5-C5-C6-O6
3	N	1	GLC	C4-C5-C6-O6
3	N	3	GLC	C4-C5-C6-O6
3	N	3	GLC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	500/529 (94%)	-1.53	0 100 100	51, 78, 114, 138	0
1	C	500/529 (94%)	-1.51	0 100 100	49, 69, 101, 121	0
2	B	30/42 (71%)	-1.61	0 100 100	75, 95, 118, 127	0
2	D	31/42 (73%)	-1.58	0 100 100	68, 81, 108, 117	0
All	All	1061/1142 (92%)	-1.52	0 100 100	49, 74, 110, 138	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GLC	M	1	12/12	-	-	56,62,70,82	0
3	GLC	M	2	11/12	-	-	57,62,68,70	0
3	GLC	M	3	11/12	-	-	60,76,93,94	0
3	GLC	M	4	11/12	-	-	94,111,121,135	0
3	GLC	N	1	12/12	-	-	51,60,67,71	0
3	GLC	N	2	11/12	-	-	49,53,59,62	0
3	GLC	N	3	11/12	-	-	55,65,72,80	0
3	GLC	N	4	11/12	-	-	88,92,99,102	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain M:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain N:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.