

Full wwPDB X-ray Structure Validation Report (i)

Dec 15, 2025 – 02:19 PM EST

PDB ID : 9ZNJ / pdb 00009znj

Title : Crystal Structure of 6,7-dimethyl-8-ribityllumazine synthase from Bordetella

pertussis in complex with 6,7-dimethyl-8-(1'-D-ribityl) lumazine

Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)

Deposited on : 2025-12-12

Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0

Mogul : 2022.3.0, CSD as543be (2022)

 $\begin{array}{ccc} \text{Xtriage (Phenix)} & : & 2.0 \\ \text{EDS} & : & 3.0 \end{array}$

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.010 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.47

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	164625	1881 (2.16-2.16)
Clashscore	180529	2047 (2.16-2.16)
Ramachandran outliers	177936	2027 (2.16-2.16)
Sidechain outliers	177891	2026 (2.16-2.16)
RSRZ outliers	164620	1882 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chair	1
			4%	
1	A	163	81%	13% 6%
	_		15%	
1	В	163	85%	8% • 7%
			6%	
1	С	163	87%	6% 7%
			9%	
1	D	163	88%	6% 6%
			8%	
1	Е	163	89%	• 7%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 5624 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 6,7-dimethyl-8-ribityllumazine synthase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	154	Total	С	N	О	S	0	0	0
1	A	104	1114	696	193	219	6	0	U	U
1	В	152	Total	С	N	О	S	0	0	0
1	Ъ	152	1072	674	182	211	5	0	U	U
1	С	152	Total	С	N	О	S	0	0	0
1		152	1096	685	190	216	5	0	0	U
1	D	153	Total	С	N	О	S	0	0	0
1	D	155	1076	675	183	212	6	0	0	U
1	Е	152	Total	С	N	О	S	0	0	0
1	Ľ	102	1072	671	184	211	6	0	U	U

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP Q7VTN4
A	-6	ALA	-	expression tag	UNP Q7VTN4
A	-5	HIS	-	expression tag	UNP Q7VTN4
A	-4	HIS	-	expression tag	UNP Q7VTN4
A	-3	HIS	-	expression tag	UNP Q7VTN4
A	-2	HIS	-	expression tag	UNP Q7VTN4
A	-1	HIS	-	expression tag	UNP Q7VTN4
A	0	HIS	-	expression tag	UNP Q7VTN4
В	-7	MET	-	initiating methionine	UNP Q7VTN4
В	-6	ALA	-	expression tag	UNP Q7VTN4
В	-5	HIS	-	expression tag	UNP Q7VTN4
В	-4	HIS	-	expression tag	UNP Q7VTN4
В	-3	HIS	-	expression tag	UNP Q7VTN4
В	-2	HIS	-	expression tag	UNP Q7VTN4
В	-1	HIS	-	expression tag	UNP Q7VTN4
В	0	HIS		expression tag	UNP Q7VTN4
С	-7	MET	=	initiating methionine	UNP Q7VTN4
С	-6	ALA	=	expression tag	UNP Q7VTN4
С	-5	HIS	-	expression tag	UNP Q7VTN4



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
С	-4	HIS	-	expression tag	UNP Q7VTN4
С	-3	HIS	-	expression tag	UNP Q7VTN4
С	-2	HIS	-	expression tag	UNP Q7VTN4
С	-1	HIS	-	expression tag	UNP Q7VTN4
С	0	HIS	-	expression tag	UNP Q7VTN4
D	-7	MET	-	initiating methionine	UNP Q7VTN4
D	-6	ALA	-	expression tag	UNP Q7VTN4
D	-5	HIS	-	expression tag	UNP Q7VTN4
D	-4	HIS	-	expression tag	UNP Q7VTN4
D	-3	HIS	-	expression tag	UNP Q7VTN4
D	-2	HIS	-	expression tag	UNP Q7VTN4
D	-1	HIS	-	expression tag	UNP Q7VTN4
D	0	HIS	-	expression tag	UNP Q7VTN4
Е	-7	MET	-	initiating methionine	UNP Q7VTN4
Е	-6	ALA	-	expression tag	UNP Q7VTN4
Е	-5	HIS	-	expression tag	UNP Q7VTN4
Е	-4	HIS	-	expression tag	UNP Q7VTN4
Е	-3	HIS	=	expression tag	UNP Q7VTN4
Е	-2	HIS	-	expression tag	UNP Q7VTN4
Е	-1	HIS	-	expression tag	UNP Q7VTN4
Е	0	HIS	=	expression tag	UNP Q7VTN4

 \bullet Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: $\mathrm{O_4P}).$



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
2	A	1	Total 5	O 4	P 1	0	0




Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total O P	0	0
	Ъ	1	5 4 1		Ŭ
2	C	1	Total O P	0	0
	C	1	5 4 1	0	U
2	D	1	Total O P	0	0
2	D	1	5 4 1		
2	E	1	Total O P	0	0
	ינו	1	5 4 1	U	0
2	E	1	Total O P	0	0
	ינו	1	5 4 1		

• Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	5	Total Cl 5 5	0	0
3	В	3	Total Cl 3 3	0	0
3	С	4	Total Cl 4 4	0	0
3	D	2	Total Cl 2 2	0	0
3	Е	2	Total Cl 2 2	0	0

• Molecule 4 is 1-deoxy-1-(6,7-dimethyl-2,4-dioxo-3,4-dihydropteridin-8(2H)-yl)-D-ribitol (CCD ID: DLZ) (formula: $C_{13}H_{18}N_4O_6$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
4	Δ	1	Total C N O	0	0	
4	Λ	1	23 13 4 6	U	0	
4	В	1	Total C N O	0	0	
4	Ъ	1	23 13 4 6	U	U	
4	\mathbf{C}	1	Total C N O	0	0	
-		1	23 13 4 6	O		
4	D	1	Total C N O	0	0	
4	D	1	23 13 4 6	U		
4	E	Г 1	Total C N O	0	0	
4	12	1	23 13 4 6		U	

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	6	Total O 6 6	0	0
5	В	4	Total O 4 4	0	0
5	С	9	Total O 9 9	0	0
5	D	5	Total O 5 5	0	0
5	Е	9	Total O 9 9	0	0




Chain E:

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 6,7-dimethyl-8-ribityllumazine synthase



89%



7%





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	135.45Å 75.42Å 94.46Å	Depositor
a, b, c, α , β , γ	90.00° 114.38° 90.00°	Depositor
Resolution (Å)	46.91 - 2.15	Depositor
Resolution (A)	46.91 - 2.15	EDS
% Data completeness	99.1 (46.91-2.15)	Depositor
(in resolution range)	99.6 (46.91-2.15)	EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.19 (at 2.16Å)	Xtriage
Refinement program	PHENIX (2.0_5904: ???)	Depositor
P. P.	0.224 , 0.261	Depositor
R, R_{free}	0.225 , 0.261	DCC
R_{free} test set	2385 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	45.0	Xtriage
Anisotropy	0.664	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 47.2	EDS
L-test for twinning ²	$ < L >=0.51, < L^2>=0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5624	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 10.36% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PO4, DLZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.21	0/1126	0.32	0/1531
1	В	0.17	0/1084	0.31	0/1479
1	С	0.21	0/1107	0.33	0/1505
1	D	0.16	0/1087	0.29	0/1482
1	Е	0.22	0/1082	0.37	0/1473
All	All	0.20	0/5486	0.33	0/7470

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1114	0	1108	17	0
1	В	1072	0	1047	10	0
1	С	1096	0	1088	10	0
1	D	1076	0	1050	9	0
1	Е	1072	0	1058	5	0
2	A	5	0	0	0	0
2	В	5	0	0	0	0
2	С	5	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	5	0	0	0	0
2	Ε	10	0	0	0	0
3	A	5	0	0	0	0
3	В	3	0	0	0	0
3	С	4	0	0	0	0
3	D	2	0	0	0	0
3	Ε	2	0	0	0	0
4	A	23	0	18	1	0
4	В	23	0	18	1	0
4	С	23	0	18	2	0
4	D	23	0	18	0	0
4	Ε	23	0	18	0	0
5	A	6	0	0	0	0
5	В	4	0	0	0	0
5	С	9	0	0	0	0
5	D	5	0	0	0	0
5	Е	9	0	0	0	0
All	All	5624	0	5441	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:51:MET:HE2	1:E:74:PHE:HZ	1.63	0.64
1:B:147:LEU:HD13	1:B:151:LEU:CD2	2.31	0.61
1:B:144:MET:HE2	1:D:60:GLU:HB3	1.85	0.59
1:A:51:MET:HE2	1:A:74:PHE:HZ	1.67	0.58
1:A:12:GLY:HA2	1:A:15:LEU:HD12	1.86	0.58
1:A:67:ARG:NH1	1:D:148:ALA:HB1	2.20	0.57
1:C:147:LEU:HD12	1:C:151:LEU:HD23	1.88	0.56
1:B:27:GLU:H	1:B:27:GLU:CD	2.13	0.56
1:E:40:LEU:HD22	1:E:45:VAL:HG21	1.88	0.56
1:A:1:MET:HG3	1:C:47:GLU:HB3	1.87	0.55
1:D:61:LEU:O	1:D:65:LEU:HD23	2.08	0.53
1:D:61:LEU:C	1:D:65:LEU:HD23	2.34	0.53
1:B:132:GLY:O	1:B:136:ASP:OD1	2.30	0.50
1:C:5:ILE:HD12	1:C:5:ILE:N	2.27	0.48
1:B:39:GLU:O	1:B:43:LEU:HD22	2.13	0.48
1:E:20:VAL:HG22	1:E:53:VAL:HG22	1.95	0.47



Continued from previous page...

A to a see 1		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap(Å)
1:C:3:PRO:O	1:C:5:ILE:HD12	2.15	0.47
1:A:133:LYS:NZ	4:C:206:DLZ:C7	2.77	0.47
1:A:16:HIS:O	1:A:17:ILE:HD13	2.14	0.47
1:A:148:ALA:HB1	1:C:67:ARG:HH12	1.79	0.47
1:A:148:ALA:HB1	1:C:67:ARG:NH1	2.30	0.47
1:B:6:LEU:C	1:B:6:LEU:HD23	2.40	0.47
1:E:20:VAL:HG22	1:E:53:VAL:CG2	2.45	0.47
1:B:53:VAL:HG21	1:E:147:LEU:HD11	1.96	0.46
1:A:47:GLU:HA	1:D:1:MET:HE3	1.97	0.46
1:A:136:ASP:O	1:A:140:VAL:HG23	2.15	0.46
1:A:50:VAL:HG21	1:D:1:MET:HE1	1.99	0.45
1:A:133:LYS:NZ	4:C:206:DLZ:H7A	2.32	0.45
1:A:38:GLU:O	1:A:42:LYS:HG2	2.17	0.44
1:C:30:GLN:O	1:C:34:GLN:HG3	2.18	0.44
1:A:83:VAL:HA	4:A:207:DLZ:O4	2.17	0.44
1:A:64:ALA:O	1:A:68:MET:HG3	2.17	0.44
1:B:152:GLU:CD	1:B:153:PRO:HD2	2.43	0.43
1:D:2:ASN:OD1	1:D:2:ASN:C	2.61	0.43
1:A:53:VAL:HG22	1:A:54:SER:N	2.33	0.43
1:C:147:LEU:HD12	1:C:147:LEU:O	2.18	0.43
4:B:205:DLZ:H7B	4:B:205:DLZ:H1'A	1.78	0.43
1:A:34:GLN:O	1:A:38:GLU:HG3	2.19	0.42
1:B:66:ALA:HA	1:B:105:ILE:HD13	2.01	0.42
1:C:147:LEU:CD1	1:C:151:LEU:HD23	2.49	0.42
1:D:66:ALA:O	1:D:70:GLU:HG2	2.19	0.42
1:C:28:ILE:O	1:C:32:GLN:HG3	2.20	0.42
1:D:6:LEU:C	1:D:6:LEU:HD23	2.46	0.41
1:B:147:LEU:HD13	1:B:147:LEU:C	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	152/163~(93%)	152 (100%)	0	0	100	100
1	В	150/163~(92%)	150 (100%)	0	0	100	100
1	С	150/163~(92%)	149 (99%)	1 (1%)	0	100	100
1	D	151/163 (93%)	150 (99%)	1 (1%)	0	100	100
1	E	150/163 (92%)	150 (100%)	0	0	100	100
All	All	753/815 (92%)	751 (100%)	2 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	110/122 (90%)	110 (100%)	0	100	100
1	В	102/122 (84%)	101 (99%)	1 (1%)	73	78
1	С	107/122 (88%)	107 (100%)	0	100	100
1	D	102/122 (84%)	102 (100%)	0	100	100
1	E	103/122 (84%)	103 (100%)	0	100	100
All	All	524/610 (86%)	523 (100%)	1 (0%)	92	95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	53	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	25	ASN
1	A	34	GLN
1	В	127	GLN



Continued from previous page...

Mol	Chain	Res	Type
1	С	16	HIS
1	С	25	ASN
1	С	90	HIS
1	С	139	GLN
1	D	16	HIS
1	D	25	ASN
1	Е	16	HIS
1	Е	127	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 27 ligands modelled in this entry, 16 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type C		Chain Res		Link	Во	ond leng	Bond angles			
MIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	Е	202	-	4,4,4	1.07	0	6,6,6	0.86	0
2	PO4	В	201	-	4,4,4	1.36	1 (25%)	6,6,6	0.69	0
2	PO4	С	201	-	4,4,4	1.46	1 (25%)	6,6,6	0.51	0
4	DLZ	D	204	-	23,24,24	1.06	1 (4%)	30,35,35	1.07	3 (10%)
2	PO4	A	201	-	4,4,4	1.53	1 (25%)	6,6,6	0.49	0



Mol	Mol Type Chain		Res	Res Link	Bo	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	tes Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	PO4	Е	201	-	4,4,4	1.60	1 (25%)	6,6,6	0.47	0	
4	DLZ	Е	205	-	23,24,24	1.17	2 (8%)	30,35,35	1.23	3 (10%)	
4	DLZ	A	207	-	23,24,24	1.12	1 (4%)	30,35,35	1.46	6 (20%)	
4	DLZ	В	205	-	23,24,24	1.13	2 (8%)	30,35,35	1.27	4 (13%)	
2	PO4	D	201	-	4,4,4	1.65	1 (25%)	6,6,6	0.50	0	
4	DLZ	С	206	-	23,24,24	1.25	2 (8%)	30,35,35	1.43	5 (16%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DLZ	D	204	-	-	3/14/14/14	0/2/2/2
4	DLZ	Е	205	-	-	3/14/14/14	0/2/2/2
4	DLZ	A	207	-	-	0/14/14/14	0/2/2/2
4	DLZ	В	205	-	-	1/14/14/14	0/2/2/2
4	DLZ	С	206	-	-	0/14/14/14	0/2/2/2

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
4	A	207	DLZ	C5A-N5	-3.53	1.31	1.39
4	С	206	DLZ	C5A-N5	-3.33	1.31	1.39
4	Е	205	DLZ	C5A-N5	-3.24	1.32	1.39
4	В	205	DLZ	C5A-N5	-3.02	1.32	1.39
2	D	201	PO4	P-O1	3.00	1.57	1.50
4	D	204	DLZ	C5A-N5	-2.99	1.32	1.39
4	С	206	DLZ	C7A-N8	-2.81	1.34	1.39
2	Е	201	PO4	P-O1	2.77	1.57	1.50
2	A	201	PO4	P-O1	2.65	1.56	1.50
2	С	201	PO4	P-O1	2.51	1.56	1.50
2	В	201	PO4	P-O1	2.50	1.56	1.50
4	Е	205	DLZ	C7A-N8	-2.26	1.35	1.39
4	В	205	DLZ	C7A-N8	-2.12	1.36	1.39

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	С	206	DLZ	C6-C5A-N5	3.85	118.20	112.45



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
4	A	207	DLZ	C7-C7A-C5A	-3.37	118.66	125.26
4	A	207	DLZ	O4-C4-N3	2.96	125.68	120.11
4	В	205	DLZ	C6-C5A-N5	2.93	116.82	112.45
4	Е	205	DLZ	C7-C7A-C5A	-2.90	119.58	125.26
4	С	206	DLZ	C7-C7A-C5A	-2.88	119.61	125.26
4	A	207	DLZ	C7-C7A-N8	2.81	121.14	116.93
4	D	204	DLZ	C6-C5A-N5	2.68	116.44	112.45
4	С	206	DLZ	C2'-C1'-N8	2.65	122.74	110.20
4	В	205	DLZ	C7-C7A-C5A	-2.64	120.09	125.26
4	Е	205	DLZ	C6-C5A-N5	2.55	116.26	112.45
4	D	204	DLZ	C7-C7A-C5A	-2.54	120.29	125.26
4	Е	205	DLZ	O4-C4-N3	2.50	124.82	120.11
4	С	206	DLZ	O4'-C4'-C3'	2.43	114.93	109.25
4	A	207	DLZ	C6-C5A-N5	2.34	115.94	112.45
4	В	205	DLZ	O4-C4-N3	2.31	124.46	120.11
4	A	207	DLZ	C2'-C1'-N8	2.27	120.92	110.20
4	С	206	DLZ	O4-C4-N3	2.18	124.22	120.11
4	A	207	DLZ	C4-C4A-N5	-2.09	115.30	118.18
4	В	205	DLZ	C7-C7A-N8	2.09	120.05	116.93
4	D	204	DLZ	O4-C4-N3	2.07	124.01	120.11

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	204	DLZ	C1'-C2'-C3'-C4'
4	Е	205	DLZ	C1'-C2'-C3'-C4'
4	Е	205	DLZ	O2'-C2'-C3'-C4'
4	D	204	DLZ	O2'-C2'-C3'-C4'
4	Е	205	DLZ	O2'-C2'-C3'-O3'
4	В	205	DLZ	O2'-C2'-C3'-C4'
4	D	204	DLZ	O2'-C2'-C3'-O3'


There are no ring outliers.

3 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	207	DLZ	1	0
4	В	205	DLZ	1	0
4	С	206	DLZ	2	0




The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	# RSRZ > 2	$OWAB(Å^2)$	Q < 0.9
1	A	154/163~(94%)	0.52	6 (3%) 44 49	48, 62, 85, 118	0
1	В	152/163~(93%)	1.09	25 (16%) 5 7	48, 80, 125, 135	0
1	С	152/163 (93%)	0.47	9 (5%) 29 35	43, 58, 87, 114	0
1	D	153/163 (93%)	1.00	14 (9%) 16 19	52, 80, 108, 123	0
1	E	152/163 (93%)	0.84	13 (8%) 18 21	45, 70, 113, 140	0
All	All	763/815 (93%)	0.78	67 (8%) 17 20	43, 69, 113, 140	0

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	153	PRO	5.7
1	D	0	HIS	5.0
1	В	153	PRO	4.6
1	В	2	ASN	4.2
1	В	4	TYR	4.1
1	Ε	4	TYR	4.0
1	С	1	MET	3.7
1	Ε	5	ILE	3.6
1	D	11	ASN	3.5
1	В	150	ALA	3.5
1	С	2	ASN	3.3
1	Ε	152	GLU	3.3
1	В	142	VAL	3.3
1	Ε	50	VAL	3.2
1	A	0	HIS	3.2
1	С	151	LEU	3.2
1	В	40	LEU	3.1
1	В	5	ILE	3.1
1	В	48	ARG	3.0
1	Ε	1	MET	3.0



Continued from previous page...

Mol	nued fron Chain	Res	Type	RSRZ
1	D	50	VAL	2.9
1	В	151	LEU	2.9
1	В	147	LEU	2.9
1	D	147	LEU	2.9
1	С	4	TYR	2.9
1	В	74	PHE	2.7
1	С	6	LEU	2.6
1	Е	7	THR	2.6
1	В	111	ILE	2.6
1	A	41	GLY	2.6
1	В	45	VAL	2.6
1	В	36	CYS	2.6
1	D	52	VAL	2.5
1	Е	72	TYR	2.5
1	D	150	ALA	2.4
1	С	73	GLU	2.4
1	D	69	ALA	2.3
1	A	136	ASP	2.3
1	Е	2	ASN	2.3
1	В	135	ALA	2.3
1	В	6	LEU	2.3
1	В	138	ALA	2.3
1	В	42	LYS	2.3
1	В	46	ASP	2.2
1	В	44	GLY	2.2
1	D	5	ILE	2.2
1	Е	6	LEU	2.2
1	С	152	GLU	2.2
1	В	148	ALA	2.2
1	D	78	ILE	2.2
1	Е	48	ARG	2.2
1	В	11	ASN	2.2
1	D	47	GLU	2.1
1	D	33	LEU	2.1
1	С	69	ALA	2.1
1	В	8	PRO	2.1
1	В	37	LEU	2.1
1	Е	14	GLY	2.1
1	Е	13	GLU	2.1
1	A	4	TYR	2.1
1	D	72	TYR	2.1
1	A	54	SER	2.1



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	6	LEU	2.1
1	Ε	43	LEU	2.1
1	В	141	ALA	2.0
1	С	5	ILE	2.0
1	D	152	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	CL	D	203	1/1	0.82	0.14	93,93,93,93	0
4	DLZ	A	207	23/23	0.84	0.14	51,62,73,76	0
4	DLZ	D	204	23/23	0.84	0.14	58,64,71,86	0
3	CL	С	202	1/1	0.86	0.15	69,69,69,69	0
3	CL	D	202	1/1	0.87	0.15	99,99,99,99	0
2	PO4	Ε	201	5/5	0.87	0.14	80,84,91,92	0
4	DLZ	E	205	23/23	0.87	0.13	42,52,63,72	0
3	CL	В	204	1/1	0.88	0.12	106,106,106,106	0
4	DLZ	С	206	23/23	0.88	0.11	38,48,55,58	0
3	CL	В	203	1/1	0.89	0.22	119,119,119,119	0
3	CL	A	204	1/1	0.90	0.20	85,85,85,85	0
3	CL	С	204	1/1	0.90	0.18	95,95,95,95	0
3	CL	A	202	1/1	0.91	0.17	80,80,80,80	0
3	CL	A	206	1/1	0.91	0.10	84,84,84,84	0
3	CL	С	205	1/1	0.91	0.11	95,95,95,95	0
3	CL	В	202	1/1	0.92	0.16	91,91,91,91	0
2	PO4	A	201	5/5	0.93	0.09	50,55,60,72	0
3	CL	A	205	1/1	0.93	0.21	107,107,107,107	0




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\AA^2)$	Q<0.9
4	DLZ	В	205	23/23	0.93	0.10	47,59,68,73	0
3	CL	Е	204	1/1	0.94	0.08	89,89,89,89	0
3	CL	A	203	1/1	0.94	0.10	78,78,78,78	0
2	PO4	Е	202	5/5	0.94	0.10	43,44,66,72	0
2	PO4	В	201	5/5	0.95	0.10	45,45,55,55	0
3	CL	С	203	1/1	0.95	0.14	92,92,92,92	0
3	CL	Е	203	1/1	0.95	0.16	87,87,87,87	0
2	PO4	D	201	5/5	0.96	0.08	46,51,57,62	0
2	PO4	С	201	5/5	0.96	0.08	46,47,55,73	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





















6.5 Other polymers (i)

There are no such residues in this entry.

